



Performance and stress levels of horses in a L1 dressage competition compared for a crossunder bitless bridle and a bridle containing a snaffle bit

Bachelor Thesis



Ananja Stehouwer
July 2014

*Performance and stress levels of horses in a L1
dressage competition compared for a crossunder bitless
bridle and a bridle containing a snaffle bit*

Name: Ananja Stehouwer
Student number: 920130001
Study: Equine, Leisure and Sports
Internal supervisor: Sylvie Beekman
External supervisor: Madeleine Calkoen, Nederlandse Vereniging Bitloos Paardrijden
Time: February 2014 - July 2014
Place: Wageningen, the Netherlands

Table of Contents

Summary	7
Samenvatting.....	8
1. Introduction	9
2. Literature review	11
2.1 Effects of different bits and bridles	11
2.2 Application of rein aids	13
2.3 Riding issues associated with bit and bridle	14
2.4 Stress in horses	15
2.5 Relationship between bridle type, performance and stress levels in dressage tests	17
3. Material and methods	18
3.1 Animals	18
3.2 Experimental design	19
3.3 Experimental procedures.....	21
4. Results	24
4.1 Salivary cortisol concentration.....	25
4.2 Heart rate frequency.....	26
4.3 Heart rate variability	26
4.4 Behaviour	27
4.5 Performance.....	28
4.6 Straightness, relaxation and bringing the horse to the hand.....	29
4.7 Rideability	30
4.5 Overview	31
5. Discussion	32
5.1 Discussion.....	32
5.2 Limitations.....	33

6. Conclusion	34
7. Recommendations	35
8. References	36
9. Annex	40
9.1 Salivary cortisol detection method	40
9.2 Test L1 number 7.....	42
9.3 Test L1 number 7 bitless.....	45
9.4 Questionnaire 1.....	48
9.5 Questionnaire 2.....	49
9.6 Heart rate per exercise.....	50
9.7 Mean beat-to-beat intervals	51
9.8 Consultation	52
9.9 Reflection on the plan of action.....	54

Summary

Since 2010 the KNHS, the Dutch equestrian association, has allowed horse-rider combinations to use a bitless bridle during official dressage tests. The crossunder bitless bridle is one of the most commonly used bridles during these tests. Research has shown that horses wearing a crossunder bridle exhibit lower heart rates compared to horses wearing a bitted bridle but conflicting results have been found concerning heart rate variability and the frequency of stress related behaviours. No performance related comparison between the crossunder bridle and bit has yet been conducted in the context of a dressage competition. Therefore the aim of this research was to compare performance and stress levels of horses with and without bit. Seven horses, Dutch Warmblood and Andalusian, performed a dressage test at the Dutch L1 level. The test was performed using a crossover design. Every horse performed once with its own bridle containing a snaffle (test 1) and once with a crossunder bridle the horses it been trained with 3 to 5 times before (test 2). Performance was judged by two official judges. A heart rate monitor was installed during the test and salivary cortisol concentration was measured with the use of Salivette rolls. A significant lower cortisol concentration of 6.6 nmol/L (± 1.845 nmol/L) was found for the crossunder bitless bridle compared to the snaffle with 8.41 nmol/L (± 3.017 nmol/L, $P=0.044$). No significant differences were found for mean heart rate ($P=0.744$), heart rate variability ($P=0.128$) and dressage test performance ($P=0.673$). Mouth opening was exhibited significantly more by the horses when wearing a snaffle ($P=0.020$). Independent of the rider's preference for the bridle with snaffle bit or the crossunder bitless bridle, horses were least stressed when riding with the crossunder bridle. It is recommended to riders to take a bitless bridle into consideration as it improves horse welfare and does not interfere with performance.

Samenvatting

Sinds 2010 staat de KNHS (Koninklijke Nederlandse Hippische Sportfederatie) ruiters toe om met een bitloos hoofdstel te starten op dressuurwedstrijden. Een van de bitloze hoofdstellen waar veelvuldig gebruik van wordt gemaakt is het kaakgekruiste hoofdstel. Uit onderzoek is gebleken dat paarden die gereden worden met een kaakgekruist hoofdstel een lagere hartslag hebben dan paarden die met bit worden gereden. Het is echter nog niet duidelijk of er ook een verschil is in de hartslagvariatie en ander stress gerelateerd gedrag. Het doel van dit onderzoek was daarom om prestatie en stress tijdens een dressuurwedstrijd te vergelijken voor paarden met en zonder bit. Zeven paarden, van de rassen KWPN en Andalusiër, namen deel aan het onderzoek. Alle paarden liepen een dressuurproef op L1 niveau. Een crossover studie werd gedaan waarbij elk paard de test één keer met het eigen hoofdstel met een trensbit reed (test 1) en één keer met een kaakgekruist hoofdstel waar van te voren 3 tot 5 keer mee getraind was (test 2). Twee officiële juryleden waren aanwezig om de proeven te beoordelen. Gedurende de proef droeg het paard een hartslagmeter en direct na de test werd een speekselmonster genomen om de concentratie van het stresshormoon cortisol te meten. Een significant verschil in cortisol concentratie werd gevonden met een concentratie van 6.6 nmol/L (± 1.845 nmol/L) voor de proeven die met het kaakgekruiste hoofdstel werden gereden en een concentratie van 8.41 nmol/L (± 3.017 nmol/L, $P=0.044$) voor de proeven met trensbit. Er was geen verschil tussen beide proeven in gemiddelde hartslag ($P=0.744$), variatie in hartslagfrequentie ($P=0.128$) en de behaalde score ($P=0.673$). Paarden die gereden werden met trensbit, sperden hun mond vaker open dan paarden gereden met kaakgekruist hoofdstel ($P=0.020$). Ook bij paarden van ruiters die de voorkeur gaven aan het rijden met trens werd een minder stress gemeten wanneer ze met kaakgekruist hoofdstel gereden werden. Het is aan te bevelen om ruiters een bitloos hoofdstel te laten proberen omdat dit paardenwelzijn verbetert en niet van invloed is op de behaalde score tijdens een dressuurproef.

1. Introduction

This thesis aims to provide more scientific evidence in a subject that has been of increasing interest during the past years: bitless riding. Using a bit to control the horse is a common method employed by practically all people engaged in equestrian activities. However this method has been questioned increasingly in the past years and many riders have begun to believe riding with a bitless bridle is a less stressful method for the horse.

In the past bitless riding was mainly recommended when a horse had mouth problems. However, during the last decade bitless riding has been introduced and accepted by many Dutch riders from another point of view: the bitless bridle is an equivalent method that may improve horse welfare and horse-rider communication (Redactie Horses, 2009a; PaardenSport, 2007). This is mainly due to the work of Dr Cook, a veterinarian and researcher specialized in diseases of the horse's mouth, ears, nose and throat. Dr Cook realized that the mouth is one of the most sensitive anatomical parts of the horse and putting a bit in it might actually be contradictory and stressful for a flight animal. As a consequence he developed a bitless bridle, allowing the rider to steer using pressure on the jaws and nose of the horse to guide it in the right direction and control its speed.

The introduction of this bridle, called the crossunder bitless bridle, caused a lot of discussion and the first riders who tried a bitless bridle were seen as real daredevils. Many people believed that without a bit in its mouth, a horse would be unstoppable and the rider no more than a passenger on its back. People were astonished when they found out a horse without bit was still controllable and could even be steered around quite nicely (van den Enk and Tulp, 2009; Calkoen, 2007; MenSport, 2007). Bitless riding during hacks and dressage training began to gain popularity and more types of bitless bridles were introduced to the market (PaardenSport, 2007).

As a result of many riders' requests to allow bitless bridles in dressage competitions, a unique event was about to happen on the 10th of July 2009. At the grounds of the KNHS (Koninklijke Nederlandse Hippische Sportfederatie, the Dutch national equestrian association) in Ermelo, the Netherlands, a bitless test competition was organized to see whether performing a dressage test in a bitless bridle is comparable to riding with a bit. Internationally, the bitless test competition of the KNHS was the first official event known to allow riders to compete in dressage using a bitless bridle in place of a bridle with a bit. It was concluded that dressage with a bitless bridle is safe and that most bridles do not reduce welfare (KNHS, 2009).

Nevertheless it there were doubts if a bitless bridle would be sufficient to use during dressage tests. Dressage is about harmony between horse and rider and an important factor for this is the horse being 'on the bit'. To allow horses with a bitless bridle in dressage competition, this definition had to be restated into 'the light contact on the reins and the head-neck position of the horse' (KNHS, 2014a).

According to the Dutch instructor and judge Johan Hamminga a horse accepting the bit and gently chewing on it is a very different situation than riding without the presence of a bit in the mouth. He sees the comparison of bitted and bitless bridles as a comparison of apples and oranges (Redactie de Paardenkrant, 2013). Hamminga was one of the attending judges during the bitless test competition of the KNHS. During the event he saw a lot of horse and rider pairs performing with a nice head-neck posture, but without a good balance. Hamminga believes the pressure of a bitless bridle on the nose does not invite the horse to really walk in a collected way like a bit can do (Redactie Horses, 2009b).

Next to this some riders have reported problems attaining the same level of bending and bringing the horse to their hand when riding bitless (Redactie Horses, 2009a).

Another main argument used in the bit vs. bitless discussion is that the type of bridle used is not that important, because it is the rider's hands that decide whether the bridle is a friendly option. A rider can do the horse damage when riding with a rough hand on a bitless bridle as well. Still a bridle with bit and a bitless bridle are two different devices, with one mainly using pressure via a metal object in the mouth and the other using pressure on the nose and jaws. According to Madeleine Calkoen, president of the Dutch association for bitless riding (Nederlandse Vereniging Bitloos Paardrijden), there are indeed differences when riding with a bitless bridle. It can help the horse to achieve smoother gaits and a better collection and the effect of seat aids is stronger (PaardenSport, 2007).

The main goal of a dressage rider should be to work together with the a horse that is a happy athlete. Secondly it is also important to have a good performance at competitions. A bridle should be able to support these goals. Therefor the objective of this research is to compare a bridle with a snaffle bit and the crossunder bitless bridle, two types of bridles that are often used by dressage riders. Firstly it aims to discover if stress levels of a horse wearing a crossunder bitless bridle are lower compared to stress levels of a horse wearing a bridle with snaffle bit. Secondly it will investigate if there are differences in the scores horse and rider receive when performing a dressage test with these bridles. Here it is also important to investigate the scores attained for straightness, relaxation and the horse searching for the rider's hands.

2. Literature review

2.1 Effects of different bits and bridles

Features of the snaffle

The snaffle bit with jointed mouthpiece is frequently used for dressage riding. It has a direct action without leverage and thus the rein pressure executed by the rider is equal to the pressure on the bit (Bennett, 2001). A single jointed mouthpiece consists of two halves or cannons connected by the joint. A double jointed mouthpiece has an intermediate piece in between the cannons and two joints. The ends of the cannons are connected to a bit ring. Common types of bit rings are the loose-ring snaffle (O-shaped) and the eggbutt snaffle (oval shaped, Figure 1).

Figure 1 From left to right: single jointed eggbutt snaffle, single jointed loose-ring snaffle and double jointed loose-ring snaffle (Agradi,2014).



Clayton et al. (1984) made use of fluoroscopic images to get a picture of what happens in the horse's mouth when applying pressure on a snaffle bit. They demonstrated that a unilateral force applied one of the reins attached to a snaffle bit causes an inward movement of the other bit ring as well and is unable to cause an independent effect on one side of the mouth. When a bilateral force is applied, the snaffle moves caudally to the tongue. The tongue may then be pressed between bit and bone and as a consequence of this some horses pull their tongue backwards. Therefore Manfredi et al. (2010) suggest a double jointed snaffle may be preferred as it accommodates the shape of the tongue better. Next to pressure on the tongue, McGreevy and McLean (2010) also identify the bars of the mouth and the lips as sites influenced by snaffle bit pressure.

Figure 2 Fluoroscopic images by Clayton et al. (1984) of the position of the snaffle bit in the horse's mouth. Left picture shows a correctly fitted and adjust snaffle bit. Right picture shows that equal force bilateral force deepens indentation of the tongue.




Figure 6. Fluoroscopic photograph of a correctly fitted and adjusted snaffle bit showing the anatomical landmarks.

M. edge of mandible	LI. lower corner incisor
P. edge of hard palate	UC. upper canine
CT. cheek teeth	LC. lower canine
UI. upper corner incisor	J. joint of bit
	R. bit ring

Clayton et al. (1984) and Uhlig (2009) have paid attention as to whether the so called 'nut cracker' effect exists. This would happen when an equal bilateral force is applied on the snaffle bit. As a result the mouthpieces are pinching the tongue and bars. Both Clayton et al. (1984) and Uhlig (2009) observed a deepening in the indentation of the tongue, but did not find the bit to contact the hard palate when rein tension was applied (Figure 2).

The crossunder bridle

The crossunder bitless bridle has been developed by a researcher on the mouth, nose, ears and throat of the horse, as a solution to problems which may be connected to the bit (Cook, 2000). The reins of the crossunder bridle are connected to straps. The straps run through a ring on the noseband, cross under the jaw and are connected to the headpiece at the top of the bridle, the place where the throat latch would normally be attached (Figure 3). The noseband is plain and not strengthened with any material that may intensify the pressure on the nose. Sites of the head where pressure is exerted when the reins are used include the ventral mandible, the nose and the poll, and to a lesser extent the cheeks (McGreevy and McLean, 2010). Pulling on one rein will give diffuse pressure on the opposite side of the head, providing a directional aid to the horse. Bilateral rein pressure results in the head being squeezed. This makes the crossunder bridle feel more like a halter for the horse, while the rider may have the feeling of riding with a normal bridle and bit (Cook, 2000).

Figure 3 The crossunder bitless bridle as seen from the side (left picture) and below (right picture) (The Bitless Bridle, 2008).

In his comparison of the crossunder bridle with other bitless bridles Cook (2000) explains why he prefers the crossunder bridle above other types of bitless bridles. Bridles like the sidepull and the hackamore work primarily by pressure on the nose. The crossunder bridle exerts pressure on less sensitive parts of the head as well and this may be more convenient to the horse. In addition, Cook (2008) explains that the design of the crossunder bridle slightly decreases rein pressure because the jaw straps are changing direction due to the metal rings applied to the noseband. Overall the intention of the crossunder bridle design is that the application of rein aids does not depend on infliction of pain. As a result the horse may not express aversive behaviours that are observed to occur when a bit is worn (Manfredi et al., 2010; Heleski et al., 2009; Ashley et al., 2005; Clayton et al., 1984).

2.2 Application of rein aids

McGreevy et al. (2011) point out that both bitted and bitless bridles rely on the use of negative reinforcement. Negative reinforcement is defined by Chance (1993) the subtraction of a stimulus after the desired behavioural change has taken place. According to McLean (2012) dressage is a process of negatively reinforcing correct responses. The bit and bridle are one of the devices the rider can use to apply negative reinforcement by relieving rein pressure when the horse gives a correct response. The goal is to diminish the rider's intervention to avoid a stress response of the horse (McGreevy and McLean, 2010). A horse's performance will improve when it is comfortable with the amount of tension the rider applies on the reins to get the desired response. In addition, this it needs to be comfortable with the mechanical effects of this tension on the bridle and bit (Manfredi et al., 2010).

A decreased responsiveness to rein aids can result when learning theory is not applied correctly by the rider. McGreevy and McLean (2005) see constant maintenance of rein pressure as the most common cause of conflict. The horse gets used to this pressure and learns to ignore it, and thus it will appear to the rider as if the horse has a hard mouth. As a solution the rider will often try to increase the pressure used or use a bit that is more severe (Warren-Smith et al., 2007).

Forces on different bridle types

The amount of pressure applied on the horse by a piece of gear depends on how much surface it is covering. McGreevy et al. (2011) explains that when this area of contact is broader, it can cause less inconvenience and thus its removal will also be less rewarding for the horse. Pesie (2010) did a calculation on the contacting surface of several bits and bridles on a model horse head. When applying a force of 50N on the model head, the amount of surface that applies pressure on the head is larger for the crossunder bitless bridle (97.5 cm²) compared to a bridle with snaffle bit (26.4 cm²). This would indicate that if a rider would apply an equal amount of tension independent of the type of bridle, a crossunder bitless bridle will cause less inconvenience. In reality, the contacting surface for a bridle containing a snaffle will be somewhat larger as Geyer and Weishaupt (2006) mention that both bitted and bitless bridles can affect facial nerves and other sensible areas of the head.

Randle and Wright (2013) did a study on the amount of rein tension applied on a crossunder bridle by riders that were only familiar to bitted bridles. The baseline tension riders applied was significantly lower for the crossunder bitless bridle compared to a jointed eggbutt snaffle. Also the tension applied when going from a walk to halt was significantly less for the crossunder bridle compared to the jointed eggbutt. Warren-Smith et al. (2007) compared rein tension between a rope halter and a bridle with a single jointed eggbutt snaffle without noseband. They did not find a difference in the mean tension applied by riders. Their study indicates a negative correlation between the rider's experience and the amount of rein tension applied.

Uhlig (2009) discovered that rein tensions of 36.8 N and higher on a snaffle bit will compress the tongue between the bit and the lower jaw and between the teeth and the premolar teeth. However the studies by Randle and Wright (2013) and Warren-Smith et al. (2007) who studied rein tensions during normal riding activities in walk and trot only found rein tensions of below 36.8 N.

2.3 Riding issues associated with bit and bridle

When comparing skeletons of domestic horses and feral horses, van Lancker et al. (2007) found that the interdental spaces of domestic horses are more irregular. They conclude that this may be caused by repeatedly induced trauma of the bit on the bars. Tell et al. (2008) found a higher occurrence of buccal ulcers in horses that were currently ridden with bit and bridle compared to horses that were not. They attribute this to poor bit fit, as well as dental procedures that are not adapted to the individual. Engelke and Gasse (2003) found asymmetrical distances between horses' left and right jaws which indicates that every horse has a different mouth shape. Nosebands can also be a source of riding issues. The noseband may cause physiological stress responses (McGreevy et al., 2012) and restrict the horse from chewing on the bit (Kienapfel and Preuschoft, 2010).

Cook (2003) and Hockenhull and Creighton (2012) indicate a difference exists between the kind of behaviour a horse shows when ridden with a bitted bridle as compared to a bitless bridle. In a survey by Cook (2003) among 605 horse owners who switched their horse from a bridle containing a bit (varying from snaffles to leverage bits) to a crossunder bitless bridle, 97% reported positive results. The main behaviours that were eliminated when riding bitless were expression of fear, difficulties slowing or stopping the horse, argumentative or aggressive behaviour, head shaking, teeth grinding, difficulties in bridling the horse, being above the bit and a lack of self-carriage. In total Cook identified 58 behaviours as being adverse effects of wearing a bit. Hockenhull and Creighton (2012) carried out a

survey among 791 leisure horses from the UK to evaluate the frequency of 15 behavioural problems in relation to the type of bridle. Horses wearing a hackamore or another type of bitless bridle were found to show a reduced likelihood of resistance or slowing down problems and discomfort compared to horses wearing a snaffle. Hockenull and Creighton (2012) did not make a distinction between bitless bridles working with and without a leverage function.

2.4 Stress in horses

A stressor is defined as an external stimulus or environmental condition to which the body gives physical or mental stress responses (Stephens, 1980). These responses are the adaptations the body needs to make in its homeostasis to function well under new conditions. When exercising, a certain amount of physiological stress is a necessary and the body's adaptations are quite similar to those made during the flight and fight response. The intensity of the flight and fight response depends on how used to the amount of exercise the animal is. In relation to animal welfare, Marlin and Nankervis (2002) consider stress being extreme adjustments a horse needs to make in order to cope with adverse effects due to its environment or management.

Competition and moreover the transport towards the competition site can be a potential stressor for the horse (Becker-Birck et al., 2013a) and may affect health, welfare and performance (Peeters et al., 2013). Peeters et al. (2013) hypothesizes that the stress/performance relationship follows an inverted U-shaped curve. A horse may sometimes be on the left side of this curve (having more stress leading to enhanced performance) or on the right side (having more stress leading to reduced performance).

Cortisol

Cortisol is one of the major stress hormones and its concentration increases 2 to 3 times during exercise (Marlin and Nankervis, 2002). Increasing cortisol levels have been found in response to stressors such as competition (Cayado et al., 2006), foundation training to prepare a horse for riding (Schmidt et al., 2010a) and transport (Schmidt et al., 2010b). Becker-Birck et al. (2013) did not find a difference in cortisol level for horses lunged in a hyperflexing and a normal head and neck position.

Blood serum analysis is a common method of measuring cortisol concentration. However, Peeters et al. (2011) and Hughes et al. (2006) validate saliva sampling as a less stressful method with a strong correlation between serum and saliva cortisol concentrations. Cortisol concentration and the period during which differences in baseline values can be detected, will vary depending on stressor intensity and duration (Strzelec et al., 2011; Jimenez et al., 1998). Peeters and al. (2011) stimulated horses with adrenocorticotrophic hormone and concluded that it took 30 minutes for salivary cortisol concentration to become significantly higher than baseline values and 180 minutes to fall back again. Peak times of 10 minutes for light exercise (Hughes et al., 2010), 20 minutes for a jumping course (Peeters et al., 2013) and 30 minutes for riding school horses being exercised (Hughes et al., 2006) have been reported.

Most research on cortisol in equines points at the existence of a circadian rhythm (e.g. Becker-Birck et al, 2013b; Gianetto et al., 2012) with salivary cortisol concentrations being highest from 9 to 11 AM (Bohák et al., 2013) and declining towards the evening (Bohák et al., 2013; Erber et al., 2013). Strzelec et al. (2011) did not find any evidence for circadian rhythm in recreational horses and their control group but did find higher morning values for jumping, dressage and eventing horses.

Peeters et al. (2013) found a positive correlation between competition performance and salivary cortisol with horses getting more penalties in a jumping course exhibiting a lower cortisol concentration. Research by Cayado et al. (2006) revealed that in contrast to jumping horses, plasma cortisol concentration of dressage horses is already different to baseline value when reaching the arena prior to competition. Cortisol concentration rose to a peak until the end of the test for all levels of dressage horses that participated (moderate, intermediate and most experienced).

Heart rate and heart rate variability

Heart rate (HR) has been found to have a positive correlation to stressors such as being caught (Jeziński et al., 1999), being handled without prior training (Visser et al., 2002) and transport (Schmidt et al., 2010b). Munsters et al. (2012) let horses approach a challenging object and found a lower heart rate when a horse was ridden by a rider that had a good match with the horse compared to a horse that was ridden by a rider that did not have a good match to the horse. Warren-Smith et al. (2007) did not find a correlation between long reining activities and an elevated heart rate frequency.

Heart rate is under control of the autonomic nervous system, with the parasympathetic (vagal) nervous system operating to relax the body and the sympathetic nervous system oppositely reacting to stress situations (Ernst, 2014). An increase in heart rate may be caused by reduced vagal activity, as the vagus nerve stimulates the parasympathetic nervous system, but also by increased sympathetic activity or even simultaneous activity within both systems. Therefore heart rate variability (HRV), the variation in beat-to-beat interval (R-R or NN) is found to be a better indicator of stress than heart rate (Von Borell et al., 2007). This is, for example, shown in research by Mohr et al. (2000) who found a significant difference in HRV but not in HR for horses mentally stressed by a removal compared to a group of control horses.

The heart of an individual with a high HRV seems to have a better ability to adapt to unpredictable stimuli (Acharya et al., 2006). HRV tends to lower, thus being more invariable, when a horse is fearful (McGreevy and McLean, 2010). However this seems to be more complicated as Mohr, Witte and Vos (2000) saw a decrease in SDANN (Standard Deviation of the averages of R-R intervals in all -minute segments of the entire recording) for physical stress while mental stress increased SDANN. Erber et al. (2013) discovered a decreased SDANN during riding for mares stabled individually compared to mares in a group housing system.

Stress related behaviours

Young et al. (2012) found a positive correlation between stress and a raised tail, flared nostrils, repeatedly looking around, pawing at the ground with front legs, stopping eating, approaching and retreating a potential stressor, stereotypic behaviour, snorting and stamping of the hind feet. Rietmann et al. (2004) found head high, explosive behaviour, deviations in speed of pace, stopping, tail swishing and defecation as behaviours indicating stress in warmblood horses during a groundwork task.

In studies concerning ridden horses gaping or chomping at the bit, head lifting, unusually high or low head carriage, head and neck movements, vocalizations, extreme ear positions, mouth opening, general signs of muscle tension, tail position and movement, bucking, rearing and shying are behaviours that have been related to discomfort (Heleski et al., 2009; Hall et al., 2013). Williams and Warren-Smith (2010) observed a competition for dressage horses and concluded that conflict responses were seen in all classes from Preliminary to Grand Prix. Being above the bit and shorting and stiffening of the stride were the responses typically shown by horses competing in the lower levels. Overall tail swishing, ears

back, being above the bit and tenseness were observed most frequently.

Horses can show hyper-reactive responses like extreme flexion and tension of the neck towards confusing or strong aids by the riders' hands (McGreevy and McLean, 2005). This can result in a loss of control by the rider. McGreevy and McLean (2005) mention that high rein pressure can result in irregular rhythms and a crooked longitudinal axis.

Manfredi et al. (2010) made use of a fluoroscopic study to view oral behaviours that are not directly visible from the outside. Behaviours that were frequently displayed when rein tension was applied were caudal retraction of the tongue, bulging the tongue between the palate and the bit and lifting the bit between the premolars. Mouthing the bit also increased with the application of rein tension, but this is regarded to be a more positive behaviour indicating acceptance. Other behaviours that were seen less frequently were moving the chin towards the chest, head shaking, flapping of the lips and flicking the tongue out of the mouth. No relationship was found between degree and type of oral behaviour and type of snaffle and behavioural patterns seem to be an individual response.

2.5 Relationship between bridle type, performance and stress levels in dressage tests

A few studies have been done on the performance of horses in a dressage test, comparing the usual bridle to the crossunder bridle. Cook and Mills (2009) found horses performed better and noted an improved horse-rider communication with the crossunder bridle. Scofield and Randle (2013) did not find a significant difference in the behaviours of tail swishing, open mouth, cow kicking, hollowing of the back and pulling down. Warren-Smith and Quick (2009) compared behaviours during foundation training (consisting of getting the horse used to a bridle, long reining and mounting preparation) and found higher frequencies of chewing, mouth opening, pawing and tail swishing in horses wearing a bridle with bit compared to horses wearing a crossunder bridle. Additionally a lower heart rate and heart rate variability was seen when long reining the bitless horses (Warren-Smith and Quick, 2009). Fejsáková et al. (2013) found a lower heart rate, a higher R-R interval and a lower, but not significant, cortisol concentration for bitless bridles compared to other bridle types.

Not only the type of bridle used, but also bit type seems to have an impact on stress. Vanderhorst et al. (2013) found horses wearing a Myler bit (allowing more freedom for the tongue) having a higher R-R interval compared to horses wearing a snaffle bit during a dressage test. Frequency of vertical head-neck fluctuations reduced significantly with the Myler bit and no difference in frequency of mouth opening, off the bit positions and performance was found.

3. Materials and methods

This study aims to draw a comparison between the bridle with snaffle bit and the crossunder bitless bridle in two ways, concerning the horse's stress levels and its performance.

The first main question is: 'Does a horse ridden in a crossunder bitless bridle experience less stress during a competition compared to a horse ridden in a bridle with a snaffle bit?' This has been analysed especially by measuring salivary cortisol concentration. Next to this data concerning heart rate frequency and heart rate were used as well as the frequency of three behaviours (tail swishing, mouth opening and head tossing) that have a relation to stress experienced by the horse when riding (Hall et al., 2013; Williams and Warren-Smith, 2010; Heleski et al, 2009).

The second main question is: 'Does a horse ridden in a crossunder bitless bridle have a higher score in a dressage test compared to a horse ridden in a bridle with a snaffle bit?' To be able to compare performance with both bridles well, the experiment was conducted in the context of a dressage competition at L1 (light) level with horse and rider consequently having experience riding dressage tests.

3.1 Animals

Nine competition horses were available for the experiment. The horses were selected for having a competition level in dressage of at least L1 (novice level). All horses were ridden by their regular rider, either the owner or leaser of the horse. Before the test day two horses (2 and 8, Table 1) dropped out due to health issues. Therefore these horses were not included in any of the analyses.

The remaining group consisted of four Dutch Warmblood horses and three Andalusian horses, all from the same stable. The horses were ranging in age from 6 to 19 (mean 10.7 years, ± 5.09 years). They were fed two meals of concentrates and two meals of roughage a day and were turned out 4 to 8 hours a day. All horses were ridden 5 to 6 times per week. A detailed overview of each horse can be found in Table 1.

Table 1 Horses stabled at the riding school the study took place

Horse number	Name	Gender	Breed	Age	Competition level*	Bit type
1	Zidane	Gelding	Dutch Warmblood	10	L1	Single jointed eggbutt snaffle
2	Excluded					
3	White	Mare	Dutch Warmblood	11	Z1	Single jointed eggbutt snaffle
4	Oficial	Gelding	Andalusian	6	L1	Double jointed loose ring snaffle
5	Caletelño	Gelding	Andalusian	6	L1	Single jointed loose ring snaffle
6	Modoso	Gelding	Andalusian	7	L1	Single jointed eggbutt snaffle
7	Maybe	Mare	Dutch Warmblood	19	M1	Single jointed loose ring snaffle
8	Excluded					
9	Ramaika	Mare	Dutch Warmblood	16	M2	Double jointed loose ring snaffle

* The foundation levels of Dutch dressage competition start with B (beginner) and are followed by L1 and L2 (light), M1 and M2 (medium), Z1 and Z2 (heavy) and ZZL (very heavy, light).

3.2 Experimental design

Twenty two days prior to the test day all horses received a fitting crossunder bitless bridle. Originally this day was also used for data collection on the bridle with snaffle. However, these measurements took place in another riding arena than the other tests and this was found to have a significant effect on the data. Thus it was decided to not to use these data.

Due to the small sample size, only one test group was defined (all animals received a crossunder bridle) instead of one test group wearing a crossunder bridle and one test group only wearing their usual bridle.. A set of 3-5 training sessions was determined to be the minimum number of times to ensure horse and rider are used to the crossunder bridle (M. Calkoen, pers. communication, 2014). These training sessions took place during the twenty two days prior to the dressage tests. Except for the bridle switch, the horse was trained in the usual way. The riders were free to choose when to schedule the training sessions with crossunder bridle.

On the test day, all horses performed a dressage test with their own bridle (test 1) and a dressage test in the crossunder bitless bridle (test 2) in an indoor riding arena of 20 meters by 40 meters (Figure 4, Figure 5). The riding arena was located at the facility where the horses were stabled and all horses were familiar with it. A crossover design was used to prevent influence of the first test on the second test as much as possible (Table 2). During the performance of a test there were no more than two horses in the arena at the same time. All horses were used to be trained with multiple horses being present in the arena and this did not seem to cause extra stress.

Table 2 Crossover schedule

Time	Horse number	Horse	Bridle
08.55	5	Caleteteño	Crossunder
09.11	1	Zidane	Crossunder
09.26	9	Ramaika	Crossunder
09.36	9	Ramaika	Snaffle
09.48	3	White	Snaffle
10.03	6	Modoso	Snaffle
10.14	4	Oficial	Snaffle
10.40	1	Zidane	Snaffle
10.42	5	Caleteteño	Snaffle
11.22	7	Maybe	Crossunder
11.36	6	Modoso	Crossunder
11.47	4	Oficial	Crossunder
11.57	3	White	Crossunder
12.05	7	Maybe	Snaffle


Before the beginning of a test, the next scheduled horse entered the arena for a warm up. Warming up had a duration of about 10 minutes. After the warming up, the heart rate monitor was installed. The horse was ridden for another 1 to 3 minutes in walk and/or trot before starting the test. At the end of the test the heart rate monitor was removed and the horse was cooled down for 3 to 4 minutes. It left the riding arena before the next scheduled horse started the test (Table 3). Upon leaving the horse waited for 5 to 10 minutes either at the washing place or in the hallway (Figure 5) to have a saliva sample taken.

Table 3 Dressage test schedule

Time (minutes)	Action
-10	Warming up horse 1
0	Installment heart rate monitor horse 1
3	Continue warming up horse 1, horse 2 enters arena for warming up
6	Start dressage test horse 1
11	End dressage test horse 1
12	Removal heart rate monitor horse 1
14	Cooling down horse 1, installment heart rate monitor horse 2
17	Horse 1 leaves riding arena
22	Saliva sample horse 1

Figure 4 (Left) The indoor riding arena seen from the viewpoint of the camera (indicated with TR in Figure 5).

Figure 5 (Right) Set up of the experiment. A riding arena of 20 meters by 40 meters was used for performance of the tests. Judges were located in opposite corners of the arena. The tripod was located behind the arena door. Installment of the heart rate monitor was done in the riding arena, saliva samples were obtained at the washing area or in the hallway of the stables.



3.3 Experimental procedures

Saliva sampling

Saliva swabs were collected using cotton rolls (Salivette; Sarstedt, Etten-Leur, Netherlands). This method has been designed for use in human research and is also applied frequently in research on horses (Hall et al., 2014; Becker-Birk et al., 2013a, 2013b; Peeters et al., 2011, 2013; Schmidt et al., 2010b). Through the longitudinal axis a cotton thread was pulled as described by Hall et al. (2014) to be able to hold the roll in place at both sides while it was inserted into the horse's mouth. It was placed on the horse's tongue just in front of the bit's position and was gently chewed on by the horse for one minute. Next the cotton roll was placed in the Salivette tube. All swabs were frozen on -20°C within 12 hours after collection. A chemiluminiscent enzyme immunoassay test was used to measure the concentration of cortisol in nmol/L (Annex 1).

Heart rate monitor

To measure heart rate, a Polar Equine RS800CX heart rate monitor was used. Time-domain measures of heart rate frequency and heart rate variability were obtained. Heart rate frequency was measured in beats per minute (bpm). Heart rate variability was measured using the standard deviation of differences between successive beat-to-beat intervals (RMSSD).

The heart rate sensor and electrodes were secured to the left D-ring at the saddle. The rider was dismounted during the placement of the heart rate monitor. The electrodes were moistened with water and a lubricant. The girth behind the horse's left elbow and a place at the right side of the horse just behind the withers were moistened as well. The electrodes were put at these places with the left electrode being secured behind the girth's straps and the right electrode being secured behind the saddle. A training watch that receives information from the sensors was placed on the rider's wrist.

All tests were filmed using a Canon SX120 IS digital camera standing on a tripod. The recording of film and heart rate monitoring were started simultaneously. The recordings and time display of the camera were used as an indication of when a certain exercise in the dressage test started. All heart rate data were processed and analysed using Polar ProTrainer 5 (Polar Electro, Kempele, Finland).

Behaviour analysis

The camera was standing on a tripod just behind the arena's fence at the flattened corner in between the letters C and H. The camera filmed continually throughout the test. The tests were analysed using an ethogram on tail, head and mouth behaviours as defined by Hall et al. (2014) and Heleski et al. (2009) (Table 4).

Table 4 Ethogram

Behaviour	Definition
<u>Tail</u>	
Swinging	Tail swinging rhythmically with body movement*
Swishing	Purposeful strong lateral and dorsoventral movements of the tail beyond that of simple rhythmic swishing*
<u>Mouth opening</u>	
Mouth still	Mouth still and shut*
Mouth open	Mouth opens and tongue or teeth may be visible*
<u>Being on the bit</u>	
Head toss	Horse moves head quickly out of 'neutral' position (the place where it is positioned for most of the ride)**

* Definitions derived from Hall et al. (2014).

**Definition derived from Heleski et al. (2009).

Performance scoring

Two judges, both registered at the KNHS and certified to judge bitless riding, scored all dressage tests. The judges were located opposite of each other with one judge in the corner in between the riding arena letters K and the A and one judge in the corner between arena letters M and the C (Figure 5). One judge (judge 1) was not familiar with the horses that were performing. The other judge (judge 2) had already judged the horses when they were riding a similar dressage test a few weeks earlier.

For the dressage test ridden with bit a test made by the KNHS was used (test L1 number 7, KNHS 2014b, Annex 2). For the bitless dressage test a similar test was used (test L1 number 7, bitloos rijden, KNHS 2014b, Annex 3). The difference between the tests was the definition used for the horse searching for the rider's hands. When a bridle with bit is used, this is explained as the horse being 'on the bit'. For bitless riding it is explained as 'a light contact on the reins and the head/neck position of the horse'.

The scoring system used for the tests was slightly adapted. Usually one score is given for the horse being straight, relaxed and on the bit. For this research straightness, relaxation and being on the bit were given a separate score.

Questionnaire

Two questionnaires were designed for evaluation of some behavioural issues of the participating horses (Annex 4, Annex 5). The questionnaire was added to the objectives of this study to fully utilize the study opportunity by hearing the experience of riders with bitless riding. To compose the questionnaires, the adverse behavioural effect list conducted by Cook (2003) was used as a guideline. The first questionnaire regarded the usual behaviour of the horse when under saddle and was filled in before the test day. The second questionnaire was specifically about the number of times the riders had been riding with the crossunder bridle and was filled in at the test day.

The questionnaires contained 21 statements about behavioural issues. By giving a score from 1 to 5 on a Likert scale (Likert, 1932) the rider indicated the frequency of the behaviour with 1 implying the horse never shows this behaviour and 5 implying the horse always shows this behaviour. Half of the

questions were stated in a positive way and the other half in a negative way. In the second questionnaire the questions were sorted in a random order. The second questionnaire also contained 8 'true or false' statements concerning the rider's experience with training the horse in the crossunder bitless bridle. Next to this riders had to rate rideability of their horse with the crossunder bridle.

Statistical analysis

Salivary cortisol concentration, heart rate, heart rate variability, behaviour frequency and total performance score were tested for normal distribution with a Kolmogorov Smirnov test. Breed differences for salivary cortisol concentration, heart rate and performance score were tested using the Independent-Samples T-Test. Differences between salivary cortisol concentration, heart rate, heart rate variability and performance score of the two bridles and were tested using a Paired Sample T-Test. For each of the 21 test exercises differences in heart rate were analysed using Kolmogorov Smirnov and a Paired Sample T-Test. The questionnaire statements were tested for differences using the Wilcoxon test. Additionally a Sign Test was used to look for positive and negative differences. For all tests the level of significance was set on $p=0.05$.

4. Results

For the first main question, ‘Does a horse ridden in a crossunder bitless bridle experience less stress during a competition compared to a horse ridden in a bridle with a snaffle bit?’, an analysis of salivary cortisol concentration was conducted. 1 out of 6 samples for test 1 (snaffle bit test) and 3 out of 6 samples for test 2 (crossunder bridle) were not available for analysis as the sample size was too small to conduct the laboratory procedure (Table 5). From one horse (horse 9) no saliva samples were obtained because this horse had to perform two consecutive tests (Table 2). Heart rate frequency and heart rate variability were analysed using the average of the entire dressage test. Heart rate frequency was also analysed using the average frequency of each separate exercise of the test. The behaviours of tail swishing, mouth opening and head tossing were scored for the entire test using an ethogram (Table 4).

Table 5 Overview of salivary cortisol samples that were successfully (✓) or not successfully (-) obtained.

Horse number	Salivary cortisol analysis	
	Test 1	Test 2
1	✓	-
3	✓	✓
4	✓	-
5	✓	-
6	-	✓
7	✓	✓
9	-	-

The second main question was concerning performance: ‘Does a horse ridden in a crossunder bitless bridle have a higher score in a dressage test compared to a horse ridden in a bridle with a snaffle bit?’. For this the total score of all 21 exercises of the dressage and the rating of the criteria for a good dressage performance (numbers 22 to 31 at of the test, Annex 2 and 3) were added up. Three separate scores, those for the criteria of straightness, relaxation and the horse searching for the rider’s hand (being ‘on the bit’), have been evaluated. Next to the riders’ experience of rideability of the horse with crossunder bridle have been compared to the stress level of the horse and the total test score. For one horse (horse 9) the questionnaire was not filled in by the rider who participated during the dressage tests but by another familiar rider.

All data were tested for breed differences between the four Dutch Warmblood horses and the three Andalusian horses. A significant difference was found for the cortisol concentration of with a mean of 10.42 nmol/L for the Dutch Warmblood and a mean of 5.39 nmol/L for the Andalusian horse ($P=0.024$). No other breed differences were discovered.

4.1 Salivary cortisol concentration


With a Kolmogorov Smirnov test a normal distribution was found for the salivary cortisol concentration of test 1 ($P=0.394$) and test 2 ($P=0.388$). For test 1 an average concentration of 8.41 nmol/L (± 3.017 nmol/L) was found. Test 2 had a mean concentration of 6.6 nmol/L (± 1.845 nmol/L, Table 6). A significant difference ($P=0.044$) was found between the concentrations of test 1 and test 2.

Table 6 Cortisol concentrations found for dressage test 1 (snaffle) and 2 (crossunder bitless bridle).

	Cortisol concentration (nmol/L)	
Horse number	Test 1 (snaffle)	Test 2 (crossunder bridle)
1	10.6	-
3	11.9	8.3
4	7.92	-
5	4.66	-
6	-	6.87
7	8.77	4.64
9	-	-
Average (st. dev.)	8.14 (± 3.017)	6.6 (± 1.845)

For two horses the cortisol concentration of both test 1 and test 2 were successfully analysed. The values found were respectively 11.9 nmol/L (test 1, snaffle) and 8.3 nmol/L (test 2, crossunder bridle) for horse no. 3 and 8.77 nmol/L (test 1) and 4.64 nmol/L (test 2) for horse no. 7 (Figure 6).

Figure 6 Salivary cortisol concentrations for individual horses after dressage test 1 and dressage test 2.



4.2 Heart rate frequency

A Kolmogorov Smirnov tests showed heart rate frequency to be normally distributed for test 1 (P=0.899) and test 2 (P=0.999) as well as minimum heart rate (P=0.956 for test 1, P=0.864 for test 2) and maximum heart rate (P=0.952 for test 1, P=0.729 for test 2). Average heart rates found for the total dressage test were 103.71 bpm (± 6.157 bpm) for test 1 and 104.29 bpm (± 7.365 bpm) for test 2 (P=0.744, Table 7). Minimum heart rate for test 1 was 76.71 bpm (± 3.039 bpm) and 76.43 bpm (± 4.276 bpm) for test 2 (P=0.766). Maximum heart rate were 132.29 bpm (± 12.737 bpm) for test 1 and 130.86 bpm (± 16.926 bpm) for test 2 (P=0.716).

There was no difference in mean heart rate between the 21 parts of test 1 and test 2. Only exercise 20 indicated a difference may exist between both tests. It consisted of a turn, a trot-halt-trot transition and another turn. For this exercise an average heart rate of 109.43 bpm (± 101.96 bpm) was found for test 1 and an average heart rate of 96.14 bpm (± 7.244 bpm) was found for test 2 (P=0.167). In total there were 8 out of 21 exercises with a higher mean heart rate of test 1 compared to test 2 and 13 out of 21 exercises with a higher mean heart rate of test 2 compared to test 1 (Annex 6).

Table 7 Individual values of mean heart rate frequency and RMSSD (standard deviation of differences between successive beat-to-beat intervals) for test 1 (snaffle bit) and test 2 (crossunder bridle).

Horse number	Heart rate frequency and st. dev. (bpm)		RMSSD (ms)	
	Test 1 (snaffle)	Test 2 (crossunder bridle)	Test 1 (snaffle)	Test 2 (crossunder bridle)
1	99 (± 9.9)	99 (± 9.8)	11	12.6
3	112 (± 14.6)	109 (± 11.3)	10.6	30.1
4	93 (± 12.4)	92 (± 12.4)	26.3	32.1
5	106 (± 14.9)	103 (± 13.1)	16.9	17.2
6	106 (± 17.7)	115 (± 25.5)	24.6	46.7
7	107 (± 13.1)	105 (± 12.4)	14.2	10.1
9	103 (± 10.5)	107 (± 9.4)	8.2	9.4
Average (st. dev.)	103.71(± 6.157)	104.29 (± 7.365)	15.97 (± 7.06)	22.70 (± 13.96)

4.3 Heart rate variability

A normal distribution was found using the Komogorov Smirnov test for mean beat to beat interval (R-R) (P=0.997 for test 1, P=0.979 for test 2), average maximum R-R (P=0.829 for test 1, P=992 for test 2) and RMSDD (P=0.966 for test 1, P=0.872 for test 2). Mean beat to beat (R-R) interval was 588.86 ms (± 34.275 ms) during test 1 and 580.71 ms (± 43.119 ms) during test 2. No significant difference in mean beat-to-beat interval was found between test 1 and test 2 (P=0.484, Annex 7). The difference in the average maximum R-R interval was 899.29 ms (± 90.917 ms) for test 1 and 949.57 ms (± 62.846 ms) for test 2 (P=0.289). RMSSD (standard deviation of differences between successive beat-to-beat intervals) was 15.97 ms (± 7.06 ms) for test 1 and 22.70 ms (± 13.958 ms) for test 2 (P=0.128) (Table 7).

4.4 Behaviour

All behaviours were shown to be normally distributed using Kolmogorov Smirnov (tail swishing $P=0.333$ for test 1, $P=0.129$ for test 2; mouth opening $P=0.939$ for test 1, $P=0.167$ for test 2; head tossing $P=0.892$ for test 1, $P=0.603$ for test 2).

In total the behaviour of tail swishing was counted 149 times for test 1 (snaffle) and 154 times for test 2 (crossunder bridle). Four horses showed less tail swishing during test 2, one horse showed more tail swishing during test 2 and two horses did not show any tail swishing. Horse 4 accounted for the main part of tail swishes by showing 76.51% of the total number of tail swishes during test 1 and 88.96% of the total number of tail swishes during test 2. The average number of tail swishes for test 1 was 21.29 (± 41.65) and 22.00 (± 50.86) for test 2 ($P=0.865$) (Table 8).

Mouth opening was counted 75 times in total for test 1 and 3 times in total for test 2. During test 1 all horses showed mouth opening on average 10.71 times (± 8.73 times). During test 2 mouth opening was only seen in two horses, on average 0.43 times (± 0.79 times) (Table 8). A significant difference ($P=0.020$) was found between the tests.

The number of head tosses was equal for both tests and had a count of 55. Four horses showed a higher frequency of head tossing during test 1 and three horses showed a higher frequency of head tossing during test 2. On average head tossing was displayed 7.86 times (± 6.04 times) during test 1 and 7.86 times (± 9.86 times) during test 2 ($P=1.000$, Table 8).

Table 8 Frequency of the behaviours of tail swishing, mouth opening and head tossing counted per test and on average.

	Number of tail swishes		Number of mouth openings		Number of head tosses	
Horse number	Test 1	Test 2	Test 1	Test 2	Test 1	Test 2
1	23	11	3	0	0	3
3	6	4	16	1	3	2
4	114	137	1	0	19	0
5	2	0	11	2	9	29
6	0	0	9	0	10	6
7	0	0	27	0	6	3
9	4	2	8	0	8	10
Average (st. dev.)	21.29 (± 41.65)	22.00 (± 50.86)	10.71 (± 8.73)	0.43 (± 0.79)	7.86 (± 6.04)	7.86 (± 9.86)

4.5 Performance

Total dressage score for test 1 given by judge 1 ($P=0.635$), judge 2 ($P=0.522$) and the average total

score of the judges for test 1 ($P=0.527$) were normally distributed. Total dressage score for test 2 given by judge 1 ($P=0.506$), judge 2 ($P=0.974$) and the average total score of the judges for test 2 ($P=0.598$) were normally distributed as well. No significant difference existed between the average dressage test scores of test 1 and test 2 ($P=0.673$) (Figure 7). Judge 1 gave a mean score of 192.14 (± 9.771 points) for test 1 and a mean score of 190.43 (± 13.939 points) for test 2 ($P=0.579$). Judge 2 gave a mean score of 195.57 (± 11.473 points) for test 1 and a mean score of 200.68 (± 14.133 points) for test 2 ($P=0.252$). Ranking of the horses was comparable for both judges. Horses with a high ranking during test 1, also received a high ranking during test 2 (Table 9).

Figure 7 Error bar presenting the average scores judge 1 and 2 gave for test 1(snaffle) and test 2 (crossunder bridle) as well as the mean of the marks for both judges combined.




Table 9 Individual marks for each test as scored by judge 1 and judge 2 as well as the average mark and ranking of each combination.

	Test 1 (snaffle)	Test 2 (crossunder bridle)
--	------------------	----------------------------

Horse number	Mark and ranking judge 1	Mark and ranking judge 2	Average mark and ranking	Mark and ranking judge 1	Mark and ranking judge 2	Average mark and ranking
1	187 (5)	196 (3)	191.5 (4)	184 (4)	202 (3)	193.0 (3)
3	212 (1)	220 (1)	216.0 (1)	217 (1)	222 (1)	219.5 (1)
4	194 (2)	191 (4)	192.5 (3)	202 (2)	218 (2)	210.0 (2)
5	194 (2)	197 (2)	195.5 (2)	183 (5)	188 (6)	185.5 (6)
6	190 (4)	188 (6)	189.0 (5)	180 (6)	193 (5)	186.5 (5)
7	182 (7)	191 (4)	189.0 (5)	187 (3)	198 (4)	192.5 (4)
9	186 (6)	186 (7)	186.0 (7)	180 (6)	185 (7)	182.5 (7)
Average (st. dev.)	192.14 (±9.771)	195.57 (±11.473)	194.21 (±10.070)	190.43 (±13.939)	200.68 (±14.133)	195.64 (±13.853)

4.6 Straightness, relaxation and bringing the horse to the hand

All three criteria were found to be normally distributed for test 1 (straightness $P=0.422$, relaxation $P=0.905$, bringing the horse to the hand $P=0.868$) and test 2 (straightness $P=0.699$, relaxation $P=0.890$, bringing the horse to the hand $P=0.819$) using a Kolmogorov Smirnov test. The average mark horses received for straightness was a 6.14 for test 1 and a 6.07 for test 2 ($P=0.604$). Relaxation was judged with an average mark of 6.0 for test 1 and 6.29 for test 2 ($P=0.321$) and bringing the horse to the hand with 5.86 for test 1 and 6.0 for test 2 ($P=0.703$). Differences in marks given for test 1 and 2 by the individual judges were not significant (Table 10), although judge 2 seemed to perceive the horses performing in the crossunder bridle as being more relaxed ($P=0.094$).

Table 10 Average scores and significance between test 1 and test 2 given for straightness, relaxation and being on the bit.

Judge	Criteria	Average mark test 1 (snaffle)	Average mark test 2 (crossunder bridle)	P-value
1	Straightness	6.14 (±0.38)	6.14 (±0.38)	1.000
	Relaxation	6.00 (±0.58)	5.86 (±0.69)	0.604
	Bringing the horse to the hand	5.86 (±0.69)	5.71 (±0.95)	0.604
2	Straightness	6.14 (±0.69)	6.00 (±0.82)	0.604
	Relaxation	6.00 (±0.58)	6.71 (±1.11)	0.094
	Bringing the horse to the hand	5.86 (±1.22)	6.29 (±1.38)	0.407
Average of both judges	Straightness	6.14 (±0.48)	6.07 (±0.54)	0.604
	Relaxation	6.00 (±0.50)	6.29 (±0.81)	0.321
	Bringing the horse to the hand	5.86 (±0.90)	6.00 (±1.16)	0.703


4.7 Rideability

Three riders (of the horses 4, 6 and 9) found it harder to bring their horses in hand when riding bitless. Four riders did not experience a difference ($P=0.083$). Three other riders (of the horses 1, 3 and 5)

perceived their horses to be less quickly relaxed, again the other riders did not note a difference ($P=0.083$).

Overall when the riders were asked to compare the rideability of their horse for the crossunder bridle with the rideability of their horse for the bridle with snaffle they rated the rideability of the horse with crossunder bridle slightly worse than rideability with the snaffle bit. A mean score of $-0.29 (\pm 2.690)$ was given. In total three riders think their performance with the crossunder bridle was worse, three think it was better and one rider was neutral (Figure 8).

Figure 8 Performance with the crossunder bridle compared with the bridle with snaffle as perceived by the horses' riders. A score of -5 indicates a very bad rideability for the crossunder bridle compared to the snaffle bit, 0 rideability is rated equal and +5 rideability for the crossunder bridle is much better.



4.8 Overview

For salivary cortisol concentration only for two horses both results were available. For both these horses the lowest concentration was found for the test with crossunder bridle. The mean heart rate was

for 3 horses the lowest during test 1, for 3 horses during test 2 and for one horse the values were exactly the same. 6 horses exhibited a higher RMSSD during test 2. Overall four horses show a consistency to have the combination of a lower heart rate and a higher RMSDD during test 2.

Four horses scored higher during the dressage test with the crossunder bridle, three horses scored higher with the snaffle bit. For the variables of straightness and bringing the horse to the hand no difference is seen. For relaxation the horses show a tendency to score higher during test 2. Stress related behaviours are displayed equally between both tests except for mouth opening which is more frequent during test 1. Only one individual horse shows a lower frequency of tail swishing during test 1 and two horses show a lower frequency of head tossing during test 1.

5. Discussion

5.1 Discussion

This study aimed to discover if there is a difference in the stress level and performance of a horse wearing a bridle containing a snaffle bit compared to a crossunder bitless bridle. It was hypothesized that a crossunder bitless bridle would cause less stress in the horse and lead to an improved performance.

An indication for a reduction in stress level of horses wearing a crossunder bridle was found measuring the salivary cortisol concentration. Research by Fejsáková et al. (2013) also found the bitless bridle group inhibited the lowest salivary cortisol concentration, although not significant.

The reduction in stress level when using the crossunder bridle is indicated by a strong tendency of the heart rate variability to increase during the tests ridden with crossunder bridle. As explained by McGreevy and McLean (2010), Borell et al. (2007) and Acharya et al. (2006) this means that when performing with the crossunder bridle the participating horses had a better ability to adapt to stimuli and were less fearful. The only exception was horse no 7 which showed a decrease in RMSSD from test 1 to test 2. This horse performed with crossunder bridle first, but this test was performed after a 40 minute break (Table 1) which may have allowed the rider to perform a longer warm up for the crossunder bridle compared to the snaffle bit. According to Mohr, Witte and Vos (2000) an increase in physical stress may have been responsible for this decrease in heart rate variability.

Mean heart rates of 103.71 bpm for test 1 and 104.29 bpm for test 2 were found. This is comparable to the research of Williams et al. (2008) who found a mean heart rate of 102 bpm (± 13 bpm) when measuring on a British dressage competition of elementary level (comparable to Dutch M2 level). Fejsáková et al. (2013) did find a significant lower heart for horses wearing a bitless bridle and Warren-Smith and Quick (2009) found horses exhibiting lower heart rates when trained with the crossunder bitless bridle.

The dressage exercise with the most difference between the heart rates (exercise 20 in the protocol, Annex 2 and 3) was an exercise with transitions that followed each other relatively quickly. Some of the riders were seen using the reins in a more sawing manner in an attempt to get the horse on the bit during the halt. According to Randle and Wright (2013) riders may have put more tension on the reins during the test with the snaffle bit compared to the test with the crossunder bridle. This may have caused an increase of conflict behaviour as Christensen et al. (2011) found a positive correlation between rein tension and conflict behaviour. Christensen et al. (2011) did not find a correlation between rein tension and heart rate, so it is not clear if there is a connection between the increased heart rate and increased rein tension with the snaffle bit during this dressage exercise.

The scores obtained during the dressage test variables formed a performance-based indication of the difference between both bridles. No significant difference was seen between test 1 and test 2 performance, although the scores for the crossunder bridle were slightly higher comparable to the research of Wittermans-Arnold (2009) and judges of this research indicate some combinations did perform better with the crossunder bridle. Wittermans-Arnold (2009) describes horse-rider combinations achieving a similar ranking independent of which bridle is performed with. No influence of the bridle on ranking can be seen in this research as well. In the research of Cook and Mills (2009) all horses improved performance significantly and no clear rank order can be distinguished. Cook and Mills (2009) rated test exercises on a scale from 0 to 10 similar to the Dutch system used in this research. Still it appears that the judge of Cook and Mills (2009) actually uses the full scale from 0 to 10 while Dutch judges are mainly used to give a score between a 4 and 7. This may explain the larger difference in performance found by Cook and Mills (2009).

Similar to this research, Wittermans-Arnold (2009) also points out riders reporting difficulties bringing

the horse to the hand. According to Cook and Mills (2009), horses with the crossunder bridle were more willing and alert, however this terminology is hard to translate to the measurements of current research.

It is interesting to note that stress level and performance of the horse may not have a relation to rideability. Some horses that were showing a lower salivary cortisol concentration and decreased stress behaviours wearing the crossunder bridle were regarded by their riders to have decreased rideability. This may well be attributed to the fact that they only practiced 3-5 times during the short time frame of 22 days and felt they needed more time to get their horse at the same level of performance with the new bridle.

5.2 Limitations

With seven participants, sample size was limited. To be able to draw far reaching conclusions, a bigger sample size is preferred. As a part of the salivary cortisol samples was not successfully analysed, this especially is a topic in need of more research.

Judging remains a subjective job and this may explain why judge 2 gave significantly higher scores compared to judge 1 when performing bitless. Also this judge had already seen all combinations participate 22 days beforehand. The judges did give similar rankings for each test, however it is better to avoid such a situation.

Because two horses were riding in the arena when a dressage test was performed, the camera was not always able to fully obtain the view of the performing horse-rider combination. Therefore some behaviours may not be noted. All behaviours were counted by hand, which made it harder to decide when a behaviour was displayed intense enough to count.

It is not found in literature whether there is useful to compare mouth opening, a stress related behaviour like related to the bit, with the frequency of mouth opening wearing the crossunder bridle. To make an honest comparison, it should be clear what the stress related signs of too much pressure at the nose are. For example a horse ridden in a bitless bridle suffering from overmuch pressure may not open its mouth but instead show increased frequency of head tossing.

A guidance is existing for correct fitting of the crossunder bridle (The Bitless Bridle, 2008). However a horse may show a preference for a certain noseband height. For example for one horse the noseband was placed at a lower place compared to the other horses. A more thorough guidance of the participating riders on bridle fitting is useful to help them recognize when the horse is comfortable.

6. Conclusion

The crossunder bridle may be an important tool to lower stress levels of dressage horses. Horses performing a dressage test in a crossunder bridle, exhibited a significantly lower salivary concentration of the stress hormone cortisol compared to their performance with a bridle containing a snaffle bit. A

higher heart rate variability, indicating less fearfulness (McGreevy and McLean, 2010), was detected in 6 out of 7 horses.

Differences in heart rate were not as pronounced as the differences in heart rate variability. During one out of the 21 exercises performed, a nearly significant difference was found, representing a lower heart rate for the crossunder bridle. The exercise consisted of a trot-halt-trot transition during which some riders used the reins more strongly to bring the horse in the hand. The pressure put on a crossunder bridle may have caused less stress than the pressure on a snaffle bit.

Removal of the bit may enable a horse to be more relaxed. However this does not necessarily include improvements in consistence of head-neck position as no difference was seen in marks for being on the bit and number of head tosses.

Three riders felt rideability of their horse was reduced when riding with the crossunder bridle. This was not reflected by the horses' stress levels which were lower when competing in the crossunder bridle. For these riders their feeling of how they performed may have a stronger connection to riding technique than to the type of bridle used. One of the horses improved test score remarkably when riding with the crossunder bridle, showing that it is possible a horse can be easier to bring to the hand wearing a crossunder bridle. Ultimately, it is the softness of the rider's hands that allow the horse to achieve a connection.

7. Recommendations

Because no significant differences between performances for bridle with snaffle bit and crossunder bridle were observed and horses ridden with the crossunder bridle were even more relaxed, serious consideration should be given to replacing the bitted bridle with the bitless one. Especially for beginning riders it is a big responsibility to be careful with the horse's mouth while they are learning how to use the rein aids.

Some of the horses participating in this research were ridden with a constant rein pressure applied. This is regarded as the most common cause of conflict by McGreevy and McLean (2005). Educating riders in the correct application of learning theory and the anatomical features of dressage may result in a greater difference in performance than using a bitless bridle instead of a snaffle bit. It is important to emphasize the need of good riding education and a proper knowledge of collection. Consequently improvements in riding skills will increase performance for any bridle used.

Next to this it is recommended for riders to compare bridles with and without bit before deciding which one to use when riding dressage. The bridle of their choice may not always be the one their horse says is best.

An issue that still needs to be tackled is the difference in performance of horses participating in higher dressage classes. It is thought by some that steering on the nose may not be sufficient to gain a sufficient level of collection. Because of this riders using a bitless bridle are not yet permitted to compete alongside riders using a bit at M1 level and higher on during official competitions. It would therefore be interesting to assess if the degree of collection using a bitless bridle is different and stays within the definition of an anatomically correctly performing horse.

A long term research, taking into account different types of bitless bridles, can give a better conclusion as to what extent performance improves riding without a bit. Research by Fejsáková et al. (2013) and Hockenhull and Creighton (2012) did not distinct between bridles applying a direct force on the nose and the hackamore applying a leverage action. It is important to make this distinction (Cook, 2000) and have separate groups of horses for each different bridle.

8. References

- Acharya U.R., Joseph K.P., Kannathal N., Lim C.M., Suri J.S. (2006). *Heart rate variability: a review*. Medical and Biological Engineering and Computing 44, 1031-1051
- Agradi (2014) *Paardenbitten*. Available at: <http://www.agradi.nl/paarden/bitten/> (Accessed: 26 June 2014).
- Ashley, F.H., Waterman-Pearson, A.E., Whay, H.R. (2005) *Behavioural assessment of pain in horses*

and donkeys: application to clinical practice and future studies. *Equine Veterinary Journal* 37(6) p565-575.

Becker-Birck, M., Schmidt, A., Lasarzika, J., Aurichb, J., Möstl, E., Auricha, C. (2013a) *Cortisol release and heart rate variability in sport horses participating in equestrian competitions*. *Journal of Veterinary Behavior* 8 (2), p87-94.

Becker-Birck, M., Schmidt, A., Wulf, M., Aurich, J., von der Wense, A., Möstl, E., Berz, R., Aurich, C. (2013b) *Cortisol release, heart rate and heart rate variability, and superficial body temperature, in horses lunged either with hyperflexion of the neck or with an extended head and neck position*. *Journal of Animal Physiology and Animal Nutrition* 97 p322-330.

Bennet, D.G. (2001) *Bits and Biting: Form and Function*. Proceedings of the 47th AAEP Annual Convention, San Diego, California, USA. P130-137.

Bohák, Z., Szabó, F., Beckers, J. -F., Melo de Sousa, N., Kutasi, O, Nagy, K., Szenci, O. (2013) *Monitoring the circadian rhythm of serum and salivary cortisol concentrations in the horse*. *Domestic Animal Endocrinology* 45 p38-42.

Calkoen, M. (2007) *Bitloos is veiliger en paardvriendelijker dan je denkt*. *Vrijruiter* 22 (4) p22-25.

Cayado, P., Muñoz-Escassi, C., Domínguez, C., Manley, W., Olabarri, B., Sánchez de la Muela, M., Castejon, F., Marañón, G., Vara, E. (2006) *Hormone response to training and competition in athletic horses*. *Equine Exercise Physiology* 7, *Equine Veterinary Journal Suppl.* 36 p274-278.

Chance, P. 1993. *Learning and Behaviour*. 3rd edn. Belmont: Brooks and Cole

Christensen, J.W., Zharkikh, T.L., Antoine, A., Malmkvist, J. (2011) *Rein tension acceptance in young horses in a voluntary test situation*. *Equine Veterinary Journal* 43 (2) p223-228.

Clayton, H.M., Lee, R. (1984) *A Fluoroscopic study of the position and action of the jointed snaffle bit in the horse's mouth*. *Journal of Equine Veterinary Science* 4 (5) p193-196.

Cook, W.R. (2000) *A solution to respiratory and other problems of the horse caused by the bit*. *Pferdeheilkunde* 16 (4) p333-351.

Cook, W.R. (2003) *Bit-induced pain: a cause of fear, flight, fight and facial neuralgia in the horse*. *Pferdeheilkunde* 19 (1) p75-82.

Cook, W.R. (2008) *Metaal in de mond: Het kruislings bitloze hoofdstel*. Mensport Books vof, Lochem, p80-91.

Cook, W.R., Mills, D.S. (2009) *Preliminary study of jointed snaffle vs. crossunder bitless bridles: Quantified comparison of behaviour in four horses*. *Equine Veterinary Journal* 41 (8) p827-830.

Engelke, E., Gasse, H. (2002) *Position of different snaffle bits inside the equine oral cavity*. *Pferdeheilkunde* 18 (4) p367-372.

Engelke, E., Gasse, H. (2003) *An anatomical study of the rostral part of the equine oral cavity with respect to position and size of a snaffle bit*. *Equine Veterinary Education* 15 (3) p158-163.

Enk, van den, M., Tulp, M. (2009) *Bitloze hoofdstellen getest en besproken*. *Bit* 171 p45-53.

Erber, R., Wulf, M., Aurich, J., Rose-Meierhöfer, S., Hoffmann, G., Möstl, E., Aurich, C. (2013) *Stress response of three-year-old horse mares to changes in husbandry system during initial equestrian training*. *Journal of Equine Veterinary Science* 33(12) p1088-1094.

Ernst, G. (2014) *Heart rate variability*. 1st edn. London, United Kingdom: Springer.

Fejsáková, M., Kottferová, J., Dankulincová, Z., Haladová, E., Matos, R., Miño, I. (2013) *Some possible factors assessing horse welfare management*. *Acta Veterinaria Brno* 82 p447-451.

Geyer, H., Weishaupt, M.A. (2006) *Der Einfluss von Zügel und Gebiss auf die Bewegungen des Pferdes - Anatomisch-funktionelle Betrachtungen*. *Pferdeheilkunde* 22 (5) p597-600.

Giannetto, C., Fazio, F., Vazzana, I., Panzera, M., Piccione, G. (2012) *Comparison of cortisol and rectal temperature circadian rhythms in horses: the role of light/dark cycle and constant darkness*. *Biological Rhythm Research* 43(6) p681-687.

- Hall, C., Huws, N., White, C., Taylor, E., Owen, H., McGreevy, P. (2013) *Assessment of ridden horse behavior*. Journal of Veterinary Behavior 8 p62-73.
- Hall, C., Kay, R., Yarnell, K. (2014) *Assessing ridden horse behavior: Professional judgment and physiological measures*. Journal of Veterinary Behaviour 9 p22-29.
- Heleski, C.R., McGreevy, P.D., Kaiser, L.J., Lavagnino, M., Tans, E., Bello, N., Clayton, H.M. (2009) *Effects on behavior and rein tension on horses ridden with or without martingales and rein inserts*. The Veterinary Journal 181 p56-62.
- Hughes, T., Creighton, E., Coleman, R. (2006) *Validation of salivary cortisol as an indicator of HPA activity in horses*. Proceedings of the 40th International Congress of the ISAE, Bristol, p157.
- Hughes, T., Creighton, E., Coleman, R. (2010) *Salivary and fecal cortisol as measures of stress in horses*. Journal of Veterinary Behavior 5(1) p60. (Abstract only)
- Irvine C. H., Alexander S.L. (1994) *Factors affecting the circadian rhythm in plasma cortisol concentration in the horse*. Domestic Animal Endocrinology 11 p227-238.
- Jezierski, T., Jaworski, Z., Górecka, A. (1999) *Effects of handling on behaviour and heart rate in Konik horses: comparison of stable and forest reared youngstock*. Applied Animal Behaviour Science 62 p1-11.
- Jimenez, M., Hinchcliff, K.W., Farris, J.W. (1997) *Catecholamines and cortisol responses of horses to incremental exertion*. Veterinary Research Communications 22(2) p107-118.
- Kienapfel, K., Preuschoft, H., (2010) *Viel zu eng! Über die Verschnallung der Nasenriemen*. Pferdeheilkunde 26 (2) p178-185.
- KNHS (2009) *Bitloos dressuurwedstrijden rijden onder de loep bij KNHS*. Available at: http://bitloos.nl/downloads/publicaties/Bitloos_dressuurwedstrijden_onder_de_loep_bij_KNHS.pdf [Accessed: 1 July 2014]
- KNHS (2014a) *Wedstrijdreglement Dressuur – Wijzigingen 2014*. Available at: <https://www.knhs.nl/media/7232966/dressuur-reglementswijzigingen-2014.pdf> [Accessed: 1 July 2014]
- KNHS (2014b) *Dressuurproeven - KNHS*. Available at: <https://www.knhs.nl/kenniscentrum/dressuur/dressuurproeven/>. [Accessed 18 March 2014].
- Lancker van, S., Broek van den, W., Simoens, P. (2007) *Incidence and morphology of bone irregularities of the equine interdental spaces (bars of the mouth)*. Equine Veterinary Education 19 (2) p103-106.
- Likert, R. (1932) *A Technique for the Measurement of Attitudes*. Archives of Psychology 140 p1-55.
- Manfredi, J.M., Rosenstein, D., Lanovaz, J.L., Nauwelaerts, S., Clayton, H.M. (2010) *Fluoroscopic study of oral behaviours in response to the presence of a bit and the effects of rein tension*. Comparative Exercise Physiology 6(4) p143-148.
- Markerink, M. (2007) *Bitloos rijden met beleid*. PaardenSport 11 p2-5.
- Marlin, D., Nankervis, K. (2002) *Equine exercise physiology*. Oxford, United Kingdom: Blackwell Publishing
- McGreevy, P., McLean, A.. (2005). Behavioural problems with the ridden horse. In: Mills, D. S., McDonnell, S. M. *The domestic horse: The origins, development and management of its behaviour*. United Kingdom: Cambridge University Press. p196-211.
- McGreevy, P., McLean, A. (2010) *Equitation Science*. 1st edn. Chichester, West Sussex, United Kingdom: Wiley-Blackwell
- McGreevy, P., McLean, A., Buckley, P., McConaghy, F., McLean, C. (2011) *How riding may affect welfare: What the equine veterinarian needs to know*. Equine Veterinary Education 23(10) p531-539.
- McGreevy, P., Warren-Smith, A., Guisard, Y. (2012) *The effect of double bridles and jaw-clamping crank nosebands on temperature of eyes and facial skin of horses*. Journal of Veterinary Behavior 7

p142-148.

McLean, A. (2012) *The positive aspects of correct negative reinforcement*. Anthrozoos: A Multidisciplinary Journal of The Interactions of People & Animals 18(3) p245-254.

MenSport (2007) *Bitloos experimenteren op Horse Event*. MenSport 7 p34,35.

Mohr, E., Witte, E., Voss, B. (2000) *Heart rate variability as stress indicator*. Archiv Tierzucht 43 p171-176.

Munsters, C.B.M., Visser, E.K., van den Broek, J., Sloet van Oldruitenborgh-Oosterbaan, M. (2012) *The influence of challenging objects and horse-rider matching on heart rate, heart rate variability and behavioural score in riding horses*. The Veterinary Journal 192 p75-80.

Peeters, M., Sulon, J., Beckers, J.-F., Ledoux, D., Vandenheede, M. (2011) *Comparison between blood serum and salivary cortisol concentrations in horses using an adrenocorticotrophic hormone challenge*. Equine Veterinary Journal 43 (4) p487-493.

Peeters, M., Closson, C., Beckers, J.-F., Vandenheede, M. (2013) *Rider and Horse Salivary Cortisol Levels During Competition and Impact on Performance*. Journal of Equine Veterinary Science 33(3) p155-160.

Pesie, S. (2010) *Een vergelijking tussen hoofdstellen met en zonder bit met betrekking tot het contactoppervlak, de inwerking en biomechanica*. Wageningen, Wageningen Universiteit.

Physick-Sheard, P., Marlin, D.J., Thornhill, R., Schroter, R.C. (2000) *Frequency domain analysis of heart rate variability in horses at rest and during exercise*. Equine Veterinary Journal 32 (3) p253-262.

Randle, H., Wright H. (2013) *Rider perception of the severity of different types of bits and the bitless bridle using rein tensionometry*. Journal of Veterinary Behavior 8 e18 (Abstract only)

Redactie Horses (2009a) *Ruim 80% ruiters voor wedstrijdrijden met bitloze optoming*. Available at: http://bitloos.nl/downloads/publicaties/enquete_bitloos_van_horses_nl.pdf [Accessed: 1 July 2014]

Redactie Horses (2009b) *Bitloos niet per definitie paardvriendelijker*. Available at: <http://www.horses.nl/opinie/09-07-24-bitloos-niet-per-definitie-paardvriendelijker/> [Accessed: 20 June 2014]

Redactie de Paardenkrant (2013) *Bitloos rijden in reguliere sport verstandige keuze?* Available at: <http://www.horses.nl/infos/bitloos-rijden/> [Accessed: 19 June 2014]

Rietmann, T.R., Stuart, A.E.A., Bernasconi, P., Stauffacher, M., Auer, J.A., Weishaupt, M.A. (2004) *Assessment of mental stress in warmblood horses: heart rate variability in comparison to heart rate and selected behavioural parameters*. Applied Animal Behaviour Science 88 p121-136.

Schmidt, A., Aurich, J., Möstl, E., Müller, J., Aurich, C. (2010a) *Changes in cortisol release and heart rate and heart rate variability during the initial training of 3-year-old sport horses*. Hormones and Behavior 58 p628-636.

Schmidt, A., Biau, S., Möstl, E., Becker-Birck, M., Morillon, B., Aurich, J., Faure, J.-M., Aurich, C. (2010b) *Changes in cortisol release and heart rate variability in sport horses during long-distance road transport*. Domestic Animal Endocrinology 38 p179-189.

Schmidt, A., Möstl, E., Wehnert, C., Aurich, J., Müller, J., Aurich, C. (2010) *Cortisol release and heart rate variability in horses during road transport*. Hormones and Behavior. 57 p209-215.

Scofield, R., Randle, H. (2013) *Preliminary comparison of behaviours exhibited by horses ridden in bitted and bitless bridles*. Journal of Veterinary Behavior 8 e1-e25. (Abstract only)

Stephens, D.B. (1980) *Stress and its measurement in domestic animals. A review of behavioural and physiological studies under field and laboratory situations*. Adv. Vet. Sci. Comp. Med. 24 p179-210.

Strzelec, K., Kankofer, M., Pietrzak, S. (2011) *Cortisol concentration in the saliva of horses subjected to different kinds of exercise*. Acta Vet. Brno 80 p101-105.

Strzelec, K., Kędzierski, W., Bereznowski, A., Janczarek, I., Bocian, K., Radosz, M. (2013) *Salivary cortisol levels in horses and their riders during three-day-events*. Bull Vet Inst Pulawy 57 p237-241.

- Tell, A., Egenvall, A., Lundström, T., Wattle, O. (2008) *The prevalence of oral ulceration in Swedish horses when ridden with bit and bridle and when unridden*. The Veterinary Journal 178 p405-410.
- The Bitless Bridle (2008) *The bitless bridle: User's manual*. Wrightsville, USA.
- The Bitless Bridle (2014). *Beta Headstall*. Available at:
http://www.bitlessbridle.com/index.php?main_page=product_info&cPath=1&products_id=1.
 [Accessed: 17 June 2014]
- Uhlig, F. (2009) *Darstellung der Lage verschiedener Trensengebisse im Pferdemaul bei Einwirkung unterschiedlich starken Zügelzuges am gerittenen Pferd im Halten*. Veterinärmedizinischen Universität Wien. Available at: <http://www.vetmeduni.ac.at/fileadmin/v/hs/hochschulschriften/bakk-pferdewissen/AC07747260.pdf> [Accessed: 3 March 2014]
- Vanderhorst, K.J., Rijksen, L., Ariëns, J., Vanriet, A., Timmerman, M., Schreuder, M., Visser, K. (2013) *The effect of type of bit on welfare and performance of horses*. Journal of Veterinary Behavior 8 (2) pe22. (Abstract only)
- Visser, E.K., van Reenen, C.G., van der Werf, J.T.N., Schilder, M.B.H., Knaap, J.H., Barneveld, A., Blokhuis, H.J. (2002) *Heart rate and heart rate variability during a novel object test and a handling test in young horses*. Physiology & Behavior 76 p289-296.
- Von Borell, E., Langbein, J., Després, G., Hansen, S., Leterrier, C., Marchant-Forde, J., Marchant-Forde, R., Minero, M., Mohr, E., Prunier, A., Valance, D., Veissier, I. (2007) *Heart rate variability as a measure of autonomic regulation of cardiac activity for assessing stress and welfare in farm animals – A review*. Physiology & Behavior 92 p293-316.
- Warren-Smith, A.K., Curtis, R.A., Greetham, L., McGreevy, P.D. (2007) *Rein contact between horse and handler during specific equitation movements*. Applied Animal Behaviour Science 108 p157-169.
- Warren-Smith, A.K., Quick, J.S. (2009) *Preliminary investigations of horses' (Equus caballus) responses to different bridles during foundation training*. Journal of Veterinary Behavior 4 (4) p169-176.
- Williams, R.J., Chandler, R.E., Marlin, D.J. (2009) *Heart rates of horses during competitive dressage*. Comparative Exercise Physiology 6 pp 7-15.
- Williams, L. R., Warren-Smith, A.K. (2010) *Conflict responses exhibited by dressage horses during competition*. Journal of Veterinary Behavior: Clinical Applications and Research 5(4) p215.
- Wittermans-Arnold (2009) *Onderzoek bitloos westernrijden. Onderzoek- en adviesituaties in de groene sector*. Stoas Hogeschool, Dronten, the Netherlands.
- Young, T., Creighton, E., Smith, T., Hosie, C. (2012) *A novel scale of behavioural indicators of stress for use with domestic horses*. Applied Animal Behaviour Science 140 p33-43.

9. Annex

9.1 Salivary cortisol detection method

IMMULITE®/IMMULITE® 1000 Cortisol

English

Intended Use: For *in vitro* diagnostic use with the IMMULITE and IMMULITE 1000 Analyzers — for the quantitative measurement of cortisol (hydrocortisone, Compound F) in serum, as an aid in the clinical assessment of adrenal status.

Catalog Number: LKCO1 (100 tests), LKCO5 (500 tests)

Test Code: COR Color: Dark Pink

Summary and Explanation

Cortisol (hydrocortisone, Compound F) is the most abundant circulating steroid and the major glucocorticoid secreted by the adrenal cortex. Physiologically effective in anti-inflammatory activity and blood pressure maintenance, cortisol is also involved in gluconeogenesis, calcium absorption and the secretion of gastric acid and pepsin.

As an indicator of adrenocortical function, measurement of blood cortisol levels is useful in the differential diagnosis of Addison's and Cushing's disease, hypopituitarism, and adrenal hyperplasia and carcinoma.¹

A variety of stimulation and suppression tests — ACTH Stimulation, ACTH Reserve, Dexamethasone Suppression — can supply supportive information on adrenocortical function.^{2,3,6}

Anomalous cortisol concentrations have been shown to exist in patients with acute infections, severe pain, diabetes mellitus or heart failure, and in women either pregnant or on estrogen therapy. In addition, certain virilizing syndromes and iatrogenic conditions raise other naturally occurring steroid levels to high (nonphysiologic) concentrations that may interfere with assaying cortisol directly, unless a highly specific antiserum is used.

Principle of the Procedure

IMMULITE/IMMULITE 1000 Cortisol is a solid-phase, competitive chemiluminescent enzyme immunoassay.

Incubation Cycles: 1 x 30 minutes.

Specimen Collection

The use of an ultracentrifuge is recommended to clear lipemic samples. Hemolyzed samples may indicate mistreatment of a specimen before receipt by the laboratory; hence the results should be interpreted with caution.

Centrifuging serum samples before a complete clot forms may result in the presence of fibrin. To prevent erroneous results due to the presence of fibrin, ensure that complete clot formation has taken place prior to centrifugation of samples. Some samples, particularly those from patients receiving anticoagulant therapy, may require increased clotting time.

Blood collection tubes from different manufacturers may yield differing values, depending on materials and additives, including gel or physical barriers, clot activators and/or anticoagulants. IMMULITE/IMMULITE 1000 Cortisol has not been tested with all possible variations of tube types.

Volume Required: 10 µL serum. (Sample cup must contain at least 100 µL more than the total volume required.)

Storage: 7 days at 2–8°C or 3 months at –20°C.⁶

Warnings and Precautions

For *in vitro* diagnostic use.

Reagents: Store at 2–8°C. Dispose of in accordance with applicable laws.

Follow universal precautions, and handle all components as if capable of transmitting infectious agents. Source materials derived from human blood were tested and found nonreactive for syphilis; for antibodies to HIV 1 and 2; for hepatitis B surface antigen; and for antibodies to hepatitis C.

Sodium azide, at concentrations less than 0.1 g/dL, has been added as a preservative. On disposal, flush with large volumes of water to prevent the buildup of potentially explosive metal azides in lead and copper plumbing.

IMMULITE/IMMULITE 1000 Cortisol (PLKCO-12, 2005-04-07)

2

Chemiluminescent Substrate: Avoid contamination and exposure to direct sunlight. (See insert.)

Water: Use distilled or deionized water.

Materials Supplied

Components are a matched set. The barcode labels are needed for the assay.

Cortisol Test Units (LCO1)

Each barcode-labeled unit contains one bead coated with polyclonal rabbit anti-cortisol antibody. Stable at 2–8°C until expiration date.

LKCO1: 100 units. LKCO5: 500 units.

Allow the Test Unit bags to come to room temperature before opening. Open by cutting along the top edge, leaving the ziplock ridge intact. Reseal the bags to protect from moisture.

Cortisol Reagent Wedge (LCO2)

With barcode. 7.5 mL alkaline phosphatase (bovine calf intestine) conjugated to cortisol in buffer, with preservative. Store capped and refrigerated; stable at 2–8°C until expiration date. Recommended usage is within 30 days after opening when stored as indicated.

LKCO1: 1 wedge. LKCO5: 5 wedges.

Cortisol Adjustors (LCOL, LCOH)

Two vials (Low and High). 3 mL each of cortisol in processed human serum, with preservative. Stable at 2–8°C for 30 days after opening, or for 6 months (aliquotted) at –20°C.

LKCO1: 1 set. LKCO5: 2 sets.

Kit Components Supplied Separately

Cortisol Sample Diluent (LCO2)

For the dilution of patient samples. One vial containing 25 mL of processed cortisol-free human serum, with preservative. Stable at 2–8°C for 30 days after opening, or for 6 months (aliquotted) at –20°C.

LSUBX: Chemiluminescent Substrate

LPWS2: Probe Wash Module

LKPM: Probe Cleaning Kit

LCHKY: Sample Cup Holders (barcoded)

LSCC: Sample Cups (disposable)

LSCC: Sample Cup Caps (optional)

IMMULITE/IMMULITE 1000 Cortisol (PLKCO-12, 2005-04-07)

3

CON6: Tri-level, multi-constituent control. Also Required. Sample transfer pipets, distilled or deionized water, controls.

Assay Procedure

Note that for optimal performance, it is important to perform all routine maintenance procedures as defined in the IMMULITE or IMMULITE 1000 Operator's Manual.

See the IMMULITE or IMMULITE 1000 Operator's Manual for: preparation, setup, dilutions, adjustment, assay and quality control procedures.

Visually inspect each Test Unit for the presence of a bead before loading it onto the system.

Recommended Adjustment Interval: 2 weeks.

Quality Control Samples: Use controls or sample pools with at least two levels (low and high) of cortisol.

Expected Values

The following reference ranges have been reported in the literature for circulating cortisol levels and for the results of stimulation and suppression tests.

Diurnal Variation	a.m.: 5–25 µg/dL (138–690 nmol/L) ^{2,3,9} p.m.: Approximately half of a.m. values
ACTH Stimulation	Over twice (usually 3 to 5 times) basal values ^{2,3}
ACTH Reserve	Below basal (control) values ^{2,3}
Dexamethasone Suppression	Below basal (control) values for the screening, low-dose and high-dose tests ^{2,3}

Consider these limits as guidelines only. Each laboratory should establish its own reference ranges.

Limitations

Circulating cortisol results may be falsely elevated in samples obtained from patients being treated with prednisone or prednisone (converted to prednisolone in

vivo). Caution must therefore be exercised with cortisol determinations for patients undergoing therapy with these and structurally related synthetic corticosteroids. Heterophilic antibodies in human serum can react with the immunoglobulins included in the assay components causing interference with *in vitro* immunoassays. [See Boscato LM, Stuart MC. Heterophilic antibodies: a problem for all immunoassays. Clin Chem 1988;34:27-33.] Samples from patients routinely exposed to animals or animal serum products can demonstrate this type of interference potentially causing an anomalous result. These reagents have been formulated to minimize the risk of interference; however, potential interactions between rare sera and test components can occur. For diagnostic purposes, the results obtained from this assay should always be used in combination with the clinical examination, patient medical history, and other findings.

Performance Data

See Tables and Graphs for data representative of the assay's performance. Results are expressed in µg/dL. (Unless otherwise noted, all were generated on serum samples collected in tubes without gel barriers or clot-promoting additives.)

Conversion Factor:

µg/dL × 27.59 → nmol/L

Calibration Range: 1 – 50 µg/dL (28 – 1,380 nmol/L).

The assay is traceable to an internal standard manufactured using qualified materials and measurement procedures.

Analytical Sensitivity: 0.2 µg/dL (5.5 nmol/L).

Precision: Samples were processed in quadruplicate over the course of 5 days, 4 runs per day, for a total of 20 runs and 80 replicates. (See "Precision" table.)

Linearity: Samples were assayed under various dilutions. (See "Linearity" table for representative data.)

Recovery: Samples spiked 1 to 19 with three cortisol solutions (109, 215 and 349 µg/dL) were assayed. (See "Recovery" table for representative data.)

Specificity: The antibody is highly specific for cortisol. (See "Specificity" table.)

Although some steroids exhibit cross-reactivity, their normal physiological concentrations are low compared to cortisol, hence they will not significantly interfere in the IMMULITE Cortisol procedure. There is, however, an observed cross-reactivity of approximately 49% with prednisone which must be considered for patients undergoing this therapy. Since prednisone is converted to prednisolone *in vivo*, caution must be exercised with cortisol determinations for patients undergoing either therapy.

Bilirubin: Samples spiked with 100 and 200 mg/dL of conjugated and unconjugated bilirubin were analyzed. Bilirubin may interfere with the assay, causing elevation of values. (See "Bilirubin" table.)

Hemolysis: Presence of packed red blood cells in concentrations up to 30 µL/mL has no effect on results, within the precision of the assay.

Lipemia: Presence of triglycerides in concentrations up to 3,000 mg/dL has no effect on results, within the precision of the assay.

Method Comparison: The assay was compared to Coat-A-Count® Cortisol on 83 patient samples. (Concentration range: approximately 3.3 to 41 µg/dL. See graph.) By linear regression:

(IML) = 0.95 (CAC) + 0.32 µg/dL

$r = 0.986$

Means:
14.6 µg/dL (IMMULITE)
15.0 µg/dL (CAC)

References

- 1) Foster L, Dunn R. Single-antibody technique for radioimmunoassay of cortisol in unextracted serum or plasma. Clin Chem 1974;20:365, 2) Farmer R, Pierce C. Plasma cortisol determinations: radioimmunoassay and competitive protein binding compared. Clin Chem 1974;20:411, 3) Ruder H, et al. A radioimmunoassay for cortisol in plasma and urine. J Clin Endo Metab 1972;35:219, 4) Rutledge B, et al. Plasma Cortisol. In: Nuclear medicine *in vitro*. 1974;120, 5) Murphy B, et al. Clinical studies utilizing a new method for the serial determination of plasma corticoids. J Canad Med Assoc 1964;90:775, 6) Sparks R. Measurement of serum 11-dihydrocortisol and cortisol after meglumine. Ann Intern Med 1971;75:717, 7) Kowalski A, Paul W. A simple extraction procedure for the determination of free (unconjugated) cortisol in urine by radioimmunoassay. Clin Chem 1979;25:1152, 8)

Burtis CA, Ashwood ER, editors. Tietz textbook of clinical chemistry, 2nd ed. Philadelphia: W.B. Saunders, 1994, 9) Wilson JD, Foster DW, editors. Williams textbook of endocrinology, 7th ed. Philadelphia: W.B. Saunders, 1985, 10) National Committee for Clinical Laboratory Standards. Procedures for the collection of diagnostic blood specimens by venipuncture; approved standard, 4th ed. NCCLS Document H3-A4, Wayne, PA: NCCLS, 1996.

Technical Assistance

In the United States, contact Siemens Healthcare Diagnostics Technical Services department. Tel: 877.229.3711. Outside the United States, contact your National Distributor.

www.siemens.com/diagnostics

The Quality System of Siemens Healthcare Diagnostics Products Ltd. is certified to ISO 13485:2003.

Tables and Graphs

Precision (µg/dL)

	Within-Run ¹			Total ²	
	Mean ³	SD ⁴	CV ⁵	SD	CV
1	2.7	0.24	8.6%	0.27	10%
2	4.4	0.33	7.5%	0.37	8.4%
3	11.3	0.88	7.8%	0.87	7.7%
4	19.0	1.1	5.8%	1.2	6.3%
5	29.1	1.8	6.2%	2.2	7.6%
6	35.8	2.4	6.7%	2.6	7.3%

Linearity (µg/dL)

	Dilution ¹		Observed ² Expected ³		%O/E ⁴
	1	16 in 16 ⁵	31.9	—	—
1	8 in 16	13.3	16.0	83%	
	4 in 16	7.5	8.0	94%	
	2 in 16	3.8	4.0	95%	
	1 in 16	1.9	2.0	95%	
2	16 in 16	34.5	—	—	
	8 in 16	15.8	17.3	91%	
	4 in 16	8.6	8.6	100%	
	2 in 16	4.7	4.3	108%	
3	16 in 16	45.3	—	—	
	8 in 16	20.9	22.7	92%	
	4 in 16	11.2	11.3	99%	
	2 in 16	5.7	5.7	100%	
	1 in 16	3.1	2.8	111%	

Recovery (µg/dL)

	Solution ¹		Observed ² Expected ³		%O/E ⁴
	1	—	10.8	—	—
1	A	16.5	15.7	105%	
	B	24.8	21.0	117%	
	C	28.4	27.7	103%	
2	—	8.2	—	—	
	A	13.8	13.2	105%	
	B	20.1	18.5	109%	
3	C	25.9	25.2	103%	
	—	16.7	—	—	
A	23.5	21.3	110%		
	29.4	26.6	111%		
B	33.9	33.3	102%		

9.2 Test L1 number 7

Proef 7 Klasse L1

Uitgave 2011

Volgnr.:

Combinatienummer:

Tijdstip:

Datum:

Ruiter:

Paard:

Jury:

Rijtijd: 5.30 minuten Rijbaan: 20x40 meter		Lichtrijden, tenzij doorzitten wordt gevraagd		Cijfer	Corr.	Bemerkingen
1	A-F-B	Binnenkomen in arbeidsdraf op de linkerhand				
2	B E	Afwenden Linkerhand				
3	F-X-M	Gebroken lijn				
4	H-X-F	Van hand veranderen en enkele passen middendraf				
5	K-X-H	Gebroken lijn, daarna doorzitten				
6	Tussen H-C-M	Overgang arbeidsstap				
7	M-E	Van hand veranderen en enkele passen middenstap				
8	Tussen E-K	Overgang arbeidsdraf doorzitten				
9		Afwenden, na enkele paardlengtes mini-maal 5 meter wijken voor het linkerbeen richting B-M, daarna rechthout en lichtrijden				
10	C-X-C Tussen C-H-E	Grote volte en na enkele drafpassen paard de hals laten strekken Teugels op maat maken				
11	E-F	Van hand veranderen				
12	Voor A A	Doorzitten Afwenden, na enkele paardlengtes minimaal 5 meter wijken voor het rechterbeen richting E-H, daarna rechthout en doorzitten				
13	Tussen C en M	Arbeidsgalop rechts aanspringen				
14	B-E-B	Grote volte en enkele sprongen middengalop				
15	Tussen F-A-K	Overgang arbeidsdraf				

16	E B	Afwenden Linkerhand				
17	Tussen C en H	Arbeidsgalop links aanspringen				
18	E-B-E	Grote volte en enkele sprongen middengalop				
19	Tussen K-A-F	Overgang arbeidsdraf				
20	Voor B B Op B-X-E lijn E	Doorzitten Afwenden Halthouden Voorwaarts in arbeidsdraf, doorzitten Linkerhand				
21	A Tussen X en G	Afwenden en arbeidsstap Halthouden en groeten Bij A in stap de rijbaan verlaten				
22		Stap				
23		Draf				
24		Galop				
25		De impuls				
26		Rechtgerichtheid				
27		Ontspanning				
28		Aanleuning				
29		De houding en zit van de ruiter/amazone				
30		Rijvaardigheid en effect van de hulpen				
31		Verzorging van het geheel	2x			
		Subtotaal				

		Strafpunten				
		Totaal				

Totaal: 320 punten

Handtekening jury:

9.3 Test L1 number 7 bitless

Proef 7 Klasse L1 Bitloos

Uitgave 2011

Datum:

Volgnr.:

Ruiter:

Combinatienummer:

Paard:

Tijdstip:

Jury:

Rijtijd: 5.30 minuten Rijbaan: 20x40 meter		Lichtrijden, tenzij doorzitten wordt gevraagd		Cijfer	Corr.	Bemerkingen
1	A-F-B	Binnenkomen in arbeidsdraf op de linkerhand				
2	B E	Afwenden Linkerhand				
3	F-X-M	Gebroken lijn				
4	H-X-F	Van hand veranderen en enkele passen middendraf				
5	K-X-H	Gebroken lijn, daarna doorzitten				
6	Tussen H-C-M	Overgang arbeidsstap				
7	M-E	Van hand veranderen en enkele passen middenstap				
8	Tussen E-K	Overgang arbeidsdraf doorzitten				
9		Afwenden, na enkele paardlengtes mini-maal 5 meter wijken voor het linkerbeen richting B-M, daarna rechthout en lichtrijden				
10	C-X-C Tussen C-H-E	Grote volte en na enkele drafpassen paard de hals laten strekken Teugels op maat maken				
11	E-F	Van hand veranderen				
12	Voor A A	Doorzitten Afwenden, na enkele paardlengtes minimaal 5 meter wijken voor het rechterbeen richting E-H, daarna rechthout en doorzitten				
13	Tussen C en M	Arbeidsgalop rechts aanspringen				
14	B-E-B	Grote volte en enkele sprongen middengalop				
15	Tussen F-A-K	Overgang arbeidsdraf				

16	E B	Afwenden Linkerhand				
17	Tussen C en H	Arbeidsgalop links aanspringen				
18	E-B-E	Grote volte en enkele sprongen middengalop				
19	Tussen K-A-F	Overgang arbeidsdraf				
20	Voor B B Op B-X-E lijn E	Doorzitten Afwenden Halthouden Voorwaarts in arbeidsdraf, doorzitten Linkerhand				
21	A Tussen X en G	Afwenden en arbeidsstap Halthouden en groeten Bij A in stap de rijbaan verlaten				
22		Stap				
23		Draf				
24		Galop				
25		De impuls				
26		Rechtgerichtheid				
27		Ontspanning				
28		Aanleuning (het lichte contact op de teugels met het paard en de hoofd/halshouding van het paard)				
29		De houding en zit van de ruiter/amazone				
30		Rijvaardigheid en effect van de hulpen				
31		Verzorging van het geheel	2x			

		Subtotaal				
		Strafpunten				
		Totaal				

Totaal: 320 punten

Handtekening jury:

9.4 Questionnaire 1

9.5 Questionnaire 2

9.6 Heart rate per exercise

Exercise number	Description	Mean HR (bpm)	Standard deviation (bpm)	Mean HR (bpm)	Standard deviation (bpm)	P-value
1	A-F-B: Enter in working trot on the left lead	101.86	11.082	96.29	12.433	0.422
2	B: turn E: left hand	103.14	13.521	103.14	13.521	0.749
3	F-X-M: loop	101.71	8.939	102.29	7.365	0.832
4	H-X-F: change of hand	104.57	8.997	106.71	7.740	0.475
5	K-X-H: loop, after which sitting trot	105.14	7.151	105.86	6.914	0.751
6	Between H-C-M: transition to working walk	98.00	5.508	99.29	6.448	0.582
7	M-E: change of hand and some strides of extended walk	82.71	5.122	84.29	5.251	0.516
8	Between E-K: transition to working trot, sitting trot	81.29	3.773	82.00	6.245	0.662
9	A: turn, after some strides yield at least 5 meters for the left leg in the direction of B-M. Next go straight on in rising trot	90.71	7.610	89.71	7.825	0.802
10	C-X-C: 20 meter circle, after a few strides full extension of the neck for some strides Between C-H-E: adjust the reins	98.57	6.680	97.57	7.743	0.708
11	E-F: change hand	101.57	5.682	101.43	6.604	0.873
12	Before A: sitting trot A: turn, after some strides yield at least 5 meters for the right leg in the direction of E-H. Next go straight on in rising trot	99.86	8.071	98.57	6.579	0.426
13	Between C and M: transition to working canter on the right lead	99.14	7.647	99.43	6.214	0.912
14	B-E-B: 20 meter circle and some strides in extended canter	113.57	10.502	117.29	11.176	0.524
15	Between F-A-K: transition to working trot	121.29	6.550	122.29	14.151	0.847
16	E: turn B: left hand	118.86	11.142	119.57	17.634	0.845
17	Between C and H: transition to working canter on the left lead	115.57	10.390	116.00	19.468	0.931
18	E-B-E: 20 meter circle and some strides in extended canter	121.71	8.712	125.00	18.547	0.622
19	Between K-A-F: transition to working trot	123.86	8.840	126.57	16.288	0.521

20	Before B: sitting trot B: turn At the B-X-E line: transition to halt. Transition to working trot, sitting trot E: left hand	111.43	11.574	109.43	10.196	0.167
21	A: turn, transition to working walk Between X and G: transition to halt, greet the judge	96.14	7.244	94.86	5.640	0.412

9.7 Mean beat-to-beat intervals

Table 1 Mean length of beat-to-beat intervals for the individual horses in milliseconds.

Horse number	Mean R-R test 1	Mean R-R test 2
1	608 (± 74)	609 (± 73)
3	537 (± 83.1)	558 (± 81)
4	640 (± 97.6)	656 (± 91)
5	574 (± 96.6)	589 (± 95.6)
6	567 (± 98.7)	520 (± 123.9)
7	614 (± 90.5)	569 (± 75.7)
9	582 (± 65.9)	564 (± 63.4)

9.8. Consultation

Date: Tuesday July 14th, at 15.30

Location: Room L563 at Van Hall Larenstein, Droevendaalsesteeg 2 in Wageningen

Duration: 50-60 minutes

Public: NVBP board members and other equestrian professionals who are familiar with the subject (KNHS representative, teachers, students)

Goal: *To discuss what influence the crossunder bitless bridle can have on the performance of horses in Dutch dressage competitions.*

Materials: Invitation letters, PowerPoint presentation, A3 papers, permanent markers, handouts, pens, drinks and food

1. Introduction (8 min)

- Overview of presentation outline
- Short introduction of the topic
 - What was the aim of the research
 - Why is this relevant
 - How does the public feel about riding bitless in dressage competitions and what problems do they see

2. “Case study” (12 min)

- Movie of the horse 'Oficial' ridden with the snaffle bit (the movie consists of some parts of the dressage test). Everybody writes down 3 things they noticed while watching.
- Movie of the horse 'Oficial' ridden with the crossunder bridle. Again everybody writes down 3 things they noticed while watching
- Group discussion to analyse and compare both movies

3. Results (15 min)

- Questions can be asked immediately during this section
- Short explanation of methodology
- Highlighting the main results
- General conclusions concerning stress level and performance of the horses

4. Recommendations and brainstorm/discussion (15 min)

- Discuss several recommendations, Ananja introduces them, acts as a discussion leader and sums up the conclusions. Depending on the available time not all items will be discussed.
 - Riders should be aware that a crossunder bitless bridle may be a good alternative for the bridle with bit. What are the main points to stress to them?
 - What are the most common reasons a try-out of a crossunder bitless bridle is not successful? What to recommend a rider who has problems having the horse performing well in a bitless bridle?
 - What kind of research concerning dressage riding with a bitless bridle would we like to see in the future? Or is the existing scientific evidence clear enough?
 - How can the NVBP use this thesis' results to support their vision and spread the knowledge about bitless riding?

5. Conclusion (3 min)

- Summarize the main outcomes and conclusions of the presentation

9.9 Reflection on the plan of action

Table 11 The plan of action as proposed in the thesis' research proposal.

Week	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26
	3 Feb	10 Feb	17 Feb	24 Feb	3 Mar	10 Mar	17 Mar	24 Mar	31 Mar	7 Apr	14 Apr	21 Apr	28 Apr	5 May	12 May	19 May	26 May	2 Jun	9 Jun	16 Jun	23 Jun
Activities																					
Feedback				x		x				x				x			x		x		
Literature review	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x	x
Writing research proposal				x	x																
Adjusting research proposal						x															
Compose test rider group						x	x														
Counsel bitless horse-rider combinations										x	x	x									
Data collecting									x				x								
Data entry and analysis										x				x	x	x					
Writing report											x	x			x	x	x	x	x		
Assessment																				x	x

Delay

The most obvious thing that struck me when I reflect on my thesis and the plan of action (Table 11) is the delay I had. Handing in the research proposal and finding a location for the research went fluent. The actual delay was caused because it was hard to plan the dates on which the test days could take place. On forehand I hadn't thought about the fact that a lot of events take place at a riding school, especially when most riders are available. Due to this I also had to plan the test days closer to each other, otherwise the riders would have had more time to practise with the bridle. The second test day was on the 11th of May, which gave me only 1.5 weeks to write a draft version and about 1 month for the final version of the thesis. During the time of having to hand in the draft we decided it would be better to hand the report in for the next deadline. This was fully agreed with by the internal supervisor, although I feel I could have seen that those deadlines were too ambitious much earlier.

When I went for the deadline of the 3rd of July, I should have made a new plan of action. This would have helped me to gain a better overview of what tasks were still ahead. I now worked too long on the result section and therefore had to rush to finish the other parts of the report in time.

Independently managing a project

I am very happy that I managed to design and execute my own research project. This is a big accomplishment for me. I had to delegate tasks, make contact with a lot of new people and had a lot of responsibilities. I even managed to enjoy my work during the test days despite of everything that asked my attention and the amount of stress that gave. Of course not everything went fluent, however I now where I made mistakes and will be extra aware of preventing them in the future.

The thesis sometimes felt as a tremendous amount of work. For the literature review I tried to split up subjects and manage deadlines for those. However I still tended to get lost into details. It would have helped when I told my deadlines to a friend who could have reminded me about them. This does not mean that I was not able to finish the literature review in a good way. I even got a compliment from a person who is much more experienced with the subject than I do about the amount of relevant research articles I found and that made me very happy. So the problem is more that I should focus on the main questions first and going more into detail once I have finished them.

Delegating tasks

While doing the research I found out that I had not delegated all tasks in a good way. I did delegate in a good way, but not always to the right persons. Sometimes I saw in the end that a person I thought capable of a job, did not execute it in the way I meant it to be. I am glad that I managed to delegate, but for those tasks it would have been better if I had done them myself.

During the test days it would have been helpful to have more volunteers around. I discovered that a nice rule of thumb would be to take one person for each task and then even ask one more volunteer. That would have enabled me to more take the role of a manager instead of having to execute several tasks at the same time.

Something else to keep in mind when you are managing a big project is to make a list of involved persons. Some people asked if I could keep them up to date but sometimes they had to remind me about that. It is a nice networking opportunity if I can send updates myself.

Enquête gedrag onder het zadel

Naast de metingen die er tijdens het rijden van de proeven verricht worden, wil ik ook graag jouw mening weten over het gedrag van je paard onder het zadel. Vul de enquête het liefst nog voor de eerste proef in.

Start

www.thesistools.com

Enquête gedrag onder het zadel

1.

Wat is je naam?*

De volgende vragen gaan over algemene gegevens van je paard.

2.

Wat is de naam van je paard?*

3.

Wat is het geslacht van je paard?*

- ☐ Merrie
- ☐ Ruin
- ☐ Hengst

4.

Wat is de leeftijd van je paard?*

5.

Wat is het ras van je paard?*

6.

Wat is jullie wedstrijdniveau (dressuur)? Als je geen wedstrijden rijdt, geef dan jullie trainingsniveau.*

7.

Welk soort bit gebruik je het gewoonlijk als je dressuur rijdt?*

- ☐ Enkelgebroken watertrens
- ☐ Enkelgebroken bustrens
- ☐ Enkelgebroken D-trens
- ☐ Dubbelgebroken watertrens
- ☐ Dubbelgebroken bustrens
- ☐ Dubbelgebroken D-trens

☐ Een ander bit, namelijk:

8.

Welke soort neusriem zit er aan het hoofdstel waar je gewoonlijk dressuur mee rijdt?*

- ☐ Gecombineerde neusriem
- ☐ Lage neusriem
- ☐ Hoge neusriem
- ☐ Mexicaanse neusriem

☐ Een andere soort neusriem, namelijk:

Verder

www.thesistools.com

De volgende stellingen over het gedrag van je paard mag je beantwoorden door een score van 1 tot 5 te geven. Vul je het meest linkse bolletje in (1), dan betekend dat dat je de beschreven situatie nooit herkend bij je paard. Het meest rechter bolletje (5) staat er voor dat je de beschreven situatie altijd herkend bij je paard. De scores die je kunt geven zijn dus van links naar rechts: nooit, zelden, regelmatig, zeer regelmatig en altijd. Als je een aanvullende opmerking hebt, kun je die onder de betreffende stelling kwijt.

9.

Het gedrag van mijn paard onder het zadel is goed te voorspellen.

Nooit

☐☐☐☐☐

Altijd

10.

Eventuele opmerkingen:

11.

Mijn paard pakt het bit tussen zijn kiezen.

Nooit

☐☐☐☐☐

Altijd

12.

Eventuele opmerkingen:

13.

Mijn paard is moeilijk terug te nemen.

Nooit

☐☐☐☐☐

Altijd

14.

Eventuele opmerkingen:

15.

Mijn paard heeft een harde mond.

Nooit

☐☐☐☐☐

Altijd

16.

Eventuele opmerkingen:

17.

Mijn paard lijkt dressuur leuk te vinden.

Nooit

☐☐☐☐☐

Altijd

18.

Eventuele opmerkingen:

19.

Mijn paard speelt graag met zijn tong en/of lippen tijdens het werk.

Nooit

☐☐☐☐☐

Altijd

20.

Eventuele opmerkingen:

21.

Mijn paard is gevoelig voor de teugelhulpen.

Nooit

☐☐☐☐☐

Altijd

22.

Eventuele opmerkingen:

23.

Mijn paard gaat met tegenzin voorwaarts tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

24.

Eventuele opmerkingen:

25.

Mijn paard schudt met zijn hoofd tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

26.

Eventuele opmerkingen:

27.

Mijn paard heeft de neiging achter het bit te kruipen.

Nooit

☐☐☐☐☐

Altijd

28.

Eventuele opmerkingen:

29.

Mijn paard staat netjes stil wanneer ik opstijg.

Nooit

☐☐☐☐☐

Altijd

30.

Eventuele opmerkingen:

31.

Mijn paard is gemakkelijk nageeflijk te rijden.

Nooit

☐☐☐☐☐

Altijd

32.

Eventuele opmerkingen:

33.

Mijn paard maakt veel speeksel aan tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

34.

Eventuele opmerkingen:

35.

Mijn paard gooit zijn tong over het bit.

Nooit

☐☐☐☐☐

Altijd

36.

Eventuele opmerkingen:

37.

Mijn paard is goed geconcentreerd tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

38.

Eventuele opmerkingen:

39.

Mijn paard stapt gemakkelijk achterwaarts onder het zadel.

Nooit

☐☐☐☐☐

Altijd

40.

Eventuele opmerkingen:

41.

Mijn paard doet zijn hoofd omhoog als ik het hoofdstel om wil doen.

Nooit

☐☐☐☐☐

Altijd

42.

Eventuele opmerkingen:

43.

Nooit

Altijd

Mijn paard ontspant zich gemakkelijk tijdens het rijden.

☐☐☐☐☐

44.

Eventuele opmerkingen:

45.

Mijn paard klemt zijn tanden op elkaar als ik het bit in wil doen.

Nooit

☐☐☐☐☐

Altijd

46.

Eventuele opmerkingen:

47.

Mijn paard is vriendelijk bij het betreden van zijn stal.

Nooit

☐☐☐☐☐

Altijd

48.

Eventuele opmerkingen:

49.

Mijn paard is kopschuw.

Nooit

☐☐☐☐☐

Altijd

50.

Eventuele opmerkingen:

51.

Mijn paard schrikt snel tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

52.

Eventuele opmerkingen:

Enquête afronden

Bedankt voor het invullen!

Enquête ervaring bitloos

Deze enquête gaat over je ervaringen met het rijden met het kaakgekruste hoofdstel. Een deel van de vragen is hetzelfde als bij de eerste enquête, maar probeer alle vragen nu te beantwoorden met de keren dat je bitloos hebt gereden in gedachten.

Start

www.thesistools.com

Enquête ervaring bitloos

1.

Wat is je naam?*

De volgende vragen gaan over de training en huisvesting van je paard, dit is om te kijken of er de afgelopen periode verschillen zijn geweest tussen de deelnemende paarden.

2.

Hoeveel keer heb je met het kaakgekruste hoofdstel gereden?*

3.

Hoeveel keer per week heb je de afgelopen tijd gemiddeld gereden? Als er ook nog andere ruiters op je paard rijden, hoeveel keer per week is er de afgelopen tijd gemiddeld op je paard gereden?*

4.

Hoe vaak per dag krijgt je paard ruwvoer?*

5.

Hoe vaak per dag krijgt je paard krachtvoer?*

6.

Hoeveel uur per dag staat je paard gemiddeld op de wei/paddock? Als je paard in de week van 18 april (toen de eerste proef plaatsvond) gemiddeld een ander aantal uur per dag op de wei/paddock stond, schrijf dat dan ook op.*

Verder

De volgende vragen gaan specifiek over het rijden met het kaakgekruste hoofdstel.

7.

Mijn paard was na 1 keer rijden gewend aan het hoofdstel.*

☐ Niet waar

☐ Waar

8.

Bij mijn paard zat er veel verschil tussen het rijden met en het rijden zonder bit.*

☐ Niet waar

☐ Waar

9.

Mijn paard ging meer in mijn handen hangen.*

☐ Niet waar

☐ Waar

10.

Ik moest meer met mijn zit rijden.*

☐ Niet waar

☐ Waar

11.

Het was moeilijker om genoeg impuls te krijgen.*

☐ Niet waar

☐ Waar

12.

Mijn paard zocht mijn hand sneller op.*

☐ Niet waar

☐ Waar

13.

Ik had meer controle over mijn paard.*

☐ Niet waar

☐ Waar

14.

Mijn paard was net zo goed recht te stellen als normaal.*

☐ Niet waar

☐ Waar

15.

Wat zijn de drie grootste veranderingen (positief of negatief) die je merkte tijdens het bitloos rijden?*

16.

Hoe goed ging het rijden met het kaakgekruste hoofdstel vergeleken met je normale hoofdstel? Geef het rijden met het kaakgekruste hoofdstel een score van -5 tot +5. (-5= het rijden ging veel slechter dan normaal, 0= het rijden ging even goed als normaal, +5= het rijden ging veel beter dan normaal)

-- maak uw keuze --

Verder

www.thesistools.com

De volgende stellingen over het gedrag van je paard tijdens het rijden met het kaakgekruste hoofdstel mag je beantwoorden door een score van 1 tot 5 te geven. Vul je het meest linkse bolletje in (1), dan betekend dat dat je de beschreven situatie nooit herkend bij je paard. Het meest rechter bolletje (5) staat er voor dat je de beschreven situatie altijd herkend bij je paard. De scores die je kunt geven zijn dus van links naar rechts: nooit, zelden, regelmatig, zeer regelmatig en altijd. Als je een aanvullende opmerking hebt, kun je die onder de betreffende stelling kwijt.

17.

Mijn paard lijkt de druk van het hoofdstel goed te accepteren.

Nooit

☐☐☐☐☐

Altijd

18.

Eventuele opmerkingen:

19.

Mijn paard staat netjes stil wanneer ik opstijg.

Nooit

☐☐☐☐☐

Altijd

20.

Eventuele opmerkingen:

21.

Mijn paard heeft de neiging achter de teugel te kruipen.

Nooit

☐☐☐☐☐

Altijd

22.

Eventuele opmerkingen:

23.

Mijn paard gaat met tegenzin voorwaarts tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

24.

Eventuele opmerkingen:

25.

Mijn paard is moeilijk terug te nemen.

Nooit

☐☐☐☐☐

Altijd

26.

Eventuele opmerkingen:

27.

Mijn paard schudt met zijn hoofd tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

Mijn paard schudt met zijn hoofd tijdens het rijden.



28.

Eventuele opmerkingen:

29.

Mijn paard speelt graag met zijn tong en/of lippen tijdens het werk.

Nooit



Altijd

30.

Eventuele opmerkingen:

31.

Mijn paard is ontspannen tijdens het optomen en rijden met het kaakgekruste hoofdstel.

Nooit



Altijd

32.

Eventuele opmerkingen:

33.

Mijn paard reageert ongevoelig op de druk op zijn neus.

Nooit

☐☐☐☐☐

Altijd

34.

Eventuele opmerkingen:

35.

Mijn paard schrikt snel tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

36.

Eventuele opmerkingen:

37.

Ik heb het gevoel dat ik veel teugeldruk moet gebruiken voordat mijn paard reageert.

Nooit

☐☐☐☐☐

Altijd

38.

Eventuele opmerkingen:

39.

Mijn paard maakt veel speeksel aan tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

40.

Eventuele opmerkingen:

41.

Mijn paard stapt gemakkelijk achterwaarts onder het zadel.

Nooit

☐☐☐☐☐

Altijd

42.

Eventuele opmerkingen:

43.

Mijn paard is kopschuw.

Nooit

☐☐☐☐☐

Altijd

44.

Eventuele opmerkingen:

45.

Mijn paard ontspant zich gemakkelijk tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

46.

Eventuele opmerkingen:

47.

Mijn paard doet zijn hoofd omhoog als ik het hoofdstel om wil doen.

Nooit

☐☐☐☐☐

Altijd

48.

Eventuele opmerkingen:

49.

Mijn paard wil de controle overnemen.

Nooit

☐☐☐☐☐

Altijd

50.

Eventuele opmerkingen:

51.

Mijn paard is vriendelijk bij het betreden van zijn stal.

Nooit

☐☐☐☐☐

Altijd

52.

Eventuele opmerkingen:

53.

Het gedrag van mijn paard onder het zadel is goed te voorspellen.

Nooit

☐☐☐☐☐

Altijd

54.

Eventuele opmerkingen:

55.

Mijn paard is gemakkelijk nageeflijk te rijden.

Nooit

☐☐☐☐☐

Altijd

56.

Eventuele opmerkingen:

57.

Mijn paard is goed geconcentreerd tijdens het rijden.

Nooit

☐☐☐☐☐

Altijd

58.

Eventuele opmerkingen:

59.

Mijn paard lijkt dressuur leuk te vinden.

Nooit

☐☐☐☐☐

Altijd

60.

Eventuele opmerkingen:

Enquête afronden

Bedankt voor het invullen!

www.thesistools.com
